

**TEAR BAG IN DIRECTION OF ARROW ACROSS
PERFORATION AND DUMP INNER BAG INTO TANK**

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Baton[®]

Herbicide

ACTIVE CONSTITUENT: 800g/kg 2,4-D present as the dimethylamine salt.

GROUP II HERBICIDE

For the control of broadleaved weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, pastures, sugar cane, peanuts and non agricultural areas as per the Directions for Use Table.

IMPORTANT: READ THE ENCLOSED LEAFLET BEFORE USE

Contents: 5kg




Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



® Baton is a registered trademark
of Nufarm Australia Limited

**WATER SOLUBLE PACKAGING
FEATURING DISSOLVE-A-BAG****Open bag as follows**

Carefully open this bag at the perforation and immediately drop the inner bag into the mix tank. Do not open or handle the inner bag - moisture will cause breakage.

**STORAGE AND DISPOSAL**

Store in the closed, original container, in a dry, cool, well ventilated area away out of direct sunlight. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty outer containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia **13 11 26**).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS)

In case of emergency: Phone (1800) 033498
Ask for shift supervisor Toll free 24 hours.

CONDITIONS OF SALE

"Nufarm Australia Limited ("Nufarm") shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."


NRA Approval No.: 46314/0902

CONVERSION CHART


Amicide 625	Baton Herbicide
320mL	250g
640mL	500g
800mL	625g
960mL	750g
1.3L	1kg
1.6L	1.25kg
1.9L	1.5kg
2.25L	1.75kg
2.6L	2kg
6.4L	5kg

To convert from Amicide 625 to 800g/kg water soluble Baton multiply Amicide 625 rate by 0.781

GUIDE FOR CORRECT USE OF WATER SOLUBLE BAGS




1. Part fill tank to a level of approx. 20% of total.



2. Add required amount of Baton Water Soluble Bags.



3. Allow bags to soak for at least 3 minutes while adding water to half fill mark, then commence agitation.*



4. Complete filling and maintain agitation.

*Tank mixing of other products should be done at point 3.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Baton[®]

Herbicide

ACTIVE CONSTITUENT: 800g/kg 2,4-D present as the dimethylamine salt.

GROUP  HERBICIDE

For the control of broadleaved weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, pastures, sugar cane, peanuts and non agricultural areas as per the Directions for Use Table.

IMPORTANT: READ THE ENCLOSED LEAFLET BEFORE USE

Contains (10) measure packs
which it is illegal to sell separately.

Contents: 5kg (10x500g)
10kg (10x1kg)

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



® Baton is a registered trademark
of Nufarm Australia Limited

STORAGE AND DISPOSAL

Store in the closed, original container, in a dry, cool, well ventilated area away out of direct sunlight. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty outer containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia **13 11 26**).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS)

In case of emergency: Phone (1800) 033498
Ask for shift supervisor Toll free 24 hours.

CONDITIONS OF SALE

"Nufarm Australia Limited ('Nufarm') shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."


NRA Approval No.: 46314/0902

CONVERSION CHART


Amicide 625	Baton Herbicide
320mL	250g
640mL	500g
800mL	625g
960mL	750g
1.3L	1kg
1.6L	1.25kg
1.9L	1.5kg
2.25L	1.75kg
2.6L	2kg
6.4L	5kg

To convert from Amicide 625 to 800g/kg water soluble Baton multiply Amicide 625 rate by 0.781

GUIDE FOR CORRECT USE OF WATER SOLUBLE BAGS




1. Part fill tank to a level of approx. 20% of total.



2. Add required amount of Baton Water Soluble Bags.



3. Allow bags to soak for at least 3 minutes while adding water to half fill mark, then commence agitation.*



4. Complete filling and maintain agitation.

*Tank mixing of other products should be done at point 3.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Baton[®]

Herbicide

ACTIVE CONSTITUENT: 800g/kg 2,4-D present as the dimethylamine salt.

NOT TO BE SOLD SEPARATELY.
BEFORE USE READ ALL DIRECTIONS ON THE OUTER PACK.

WATER SOLUBLE PACKAGING: KEEP DRY

Contents: 500g
1kg

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001

NRA Approval No.: 46314/0902

® Baton is a registered trademark
of Nufarm Australia Limited



CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Baton[®]

Herbicide

ACTIVE CONSTITUENT: 800g/kg 2,4-D present as the dimethylamine salt.

GROUP	II	HERBICIDE
-------	----	-----------

For the control of broadleaved weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, pastures, sugar cane, peanuts and non agricultural areas as per the Directions for Use Table.

READ THOROUGHLY BEFORE USING.

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



® Baton is a registered trademark
of Nufarm Australia Limited

NRA Approval No.: 46314/0902

DIRECTIONS FOR USE:**RESTRAINTS:** DO NOT spray if rain seems likely within 6 hours or if strong winds prevail.

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

1. FIELD CROPS

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Wheat	Refer Weed Table	NSW, SA only	250g to 1.3kg	Apply after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller (NSW, SA only).
		Vic only	250g to 1.1kg	
		Qld only	400g to 1.1kg	Apply from tillering to boot stage. (Vic only)
		Tas only	1.1kg	Apply from mid tillering to before boot stage. (Qld only)
Barley	Refer Weed Table	NSW, SA only	250g to 1.1kg	Apply from the 5 leaf to fully tillered stage. (Tas only)
		Vic only	250g to 1.1kg	
		Qld only	400g to 1.1kg	
		Tas only	1.1kg	
Cereal Rye, Triticale	Refer Weed Table	NSW, SA only	250g to 1.1kg	
		Vic only	250g to 1.1kg	
Oats	Refer Weed Table	NSW, SA only	250 to 600g	
		Vic only	250 to 600g	
Cereals: Wheat, Oats, Barley	Cape Tulip	WA only	500g to 1kg	Apply from the 5 leaf stage up to jointing stage (Zadoks 15-33). Apply after the 6 leaf stage (Z. 16) for Cranbrook, Jacup, Aroona and Spear Wheat and Mortlock Oats to avoid possible damage. DO NOT spray if lucerne is present. WEED STAGE: 10-15cm. Docks should be sprayed before 5 leaf stage. Cape Tulip - low rate for cormils only.
	Dock, Saffron Thistle		900g	
	Indian Hedge Mustard, London Rocket, Lupin, Rapistrum, Wild Radish		600g	
	Wild Turnip		500g	
	Capeweed, Doublegee, Erodium, London Rocket, Lupin, Mustard, Rapistrum, Wild Radish, Wild Turnip		160g/ha plus 500mL/ha Nufarm Flowable Diuron (500g/L)	Apply when crop has 4-5 leaves and most weeds have germinated and are in 2-5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown medics.
Fallow, Stubble Spray prior to Direct Drilling or Sowing a) Winter Cereals b) Winter Cereals and Peanuts	Refer Weed Table	Vic only	250g to 1.1kg	Observe plant back periods given in the table on this leaflet. Can be mixed with Glean*, Siege*, Nuquat®, or Spray-Seed* where grasses are present.
		NSW & Qld only	400g to 1.3kg	For skeleton weed spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.
Millet	Refer Weed Table	NSW, SA, Vic only	400 to 900g	Spray when secondary roots have developed, when fully tillered and before heads start to form at the base of the tillers. Do not use on Panorama Millet or Panicum.
		Qld only	400 to 700g	
Sacaline, Broom Millet, Millet	Cape Tulip, Dock, Saffron Thistle, Indian Hedge Mustard, London Rocket, Lupin, Rapistrum, Radish, Wild Turnip	WA only	900g	Spray when crop is 10-30cm high and secondary roots have developed and before tasselling. Apply as direct spray to weeds.
Sugar Cane	Bindy Eye (Star Burr), Blue Top, Cobblers Pegs, Fleabanes, Jute, Leucas, Needle Burr, Spear Thistle, Water Primrose, Ipomea Vines, Convolvulus Vines	Qld only	1.4 to 2.8kg	Add 175-350mL Spraymate Activator/100L of spray mixture. Agitate well. DO NOT USE ON Q63, Q67, Q80, or Q96 VARIETIES.
	Chinese Mint, Blue Snakeweed		2.8kg	

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Peanuts	Broadleaf weeds, except Noogoora Burr. Grasses, except Mossman Burr	Qld only	1.4 or 2.8kg	LOWER RATE: Apply as BAND SPRAY as soon as possible after planting in a 55cm band. HIGHER RATE: Apply as OVERALL SPRAY after planting and before crop emergence. Some crop damage may occur if heavy rain falls between application and crop emergence.
Harvest Aid or Salvage Spray -Winter Cereals	Dessicate Broadleaf Weeds	Qld only	900g to 1.3kg	Apply after dough stage.
	Refer Weed Table	Qld & NSW only		

2. PASTURES, NON-AGRICULTURAL, RIGHTS OF WAY, INDUSTRIAL, LAWNS

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Pastures and Non-agricultural	Refer Weed Table	NSW, Qld, SA, Tas only	400g to 1.3kg	Pasture legumes including lucerne, clovers and medics may be damaged unless well protected by grasses. Spot spraying is preferred.
	Amsinckia, Docks, Bindweed, Caltrop, Flatweed, Spear Thistle, Capeweed, Doublegee, Saffron Thistle, Mustard, Wild Radish, Wild Turnip, Annual Thistles, Patersons Curse	WA only	900g	For pastures not containing legumes. Only seedling Docks, Spear Thistle and Saffron Thistle will be controlled.
	Afghan Melons	WA only	1.3kg/ha plus 1% crop oil	Spray when plants are actively young preferably before flowering or vining.
	Paddy Melons		600 to 900g	
	Prickly Saltwort (Roly Poly)		1.3kg	Spray when plants are small.
	Stinkwort		1.3 to 2.5kg plus surfactant	Best results are obtained when plants are small. Use high rate on larger plants.
	Dove Weed		2.5kg	Spray after good emergence of seedlings.
Pastures, Rights of Way & Industrial	Boxthorn, Boneseed, Hawthorn	Vic, SA only	500g/100L water	Spot Spraying. For Boneseed only, thoroughly wet plants or seedlings.
			1kg in 1.6L water	Apply to freshly cut stumps.
	Groundsel	NSW, Qld, SA only	500g/10L water	MISTING: Lightly wet plants.
			250g/100L water	HIGH VOLUME: Thoroughly wet plants.
			125g/10L water	CUT STUMP: Swab the cut stump within one hour of cutting. Apply by a pouring can or knapsack spray.
			2.3 to 3.4kg	AERIAL APPLICATION: Spray when Groundsel is actively growing.
	Lantana	NSW, Qld, SA only	250g/100L water	Use a coarse spray with sufficient pressure to penetrate canopy and wet stems as well as foliage. Spray at the end of a wet summer (March to May). Defoliation should occur but respraying of new growth will be necessary in following Autumn. Broadcast grass seed and keep stock off following Summer to allow the pasture to establish. Damage may result to pasture legumes.

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Pastures, Rights of Way & Industrial (cont.)	Mother of Millions	NSW only	300g/100L water	Hand gun and Knapsack only. A thorough coverage of leaves and plantlets is necessary. Use Spraymate Chemwet 1000 at the rate of 1mL of surfactant/1L of spray mixture.
	Noogoora Burr, Weir Vine (Ipomea)	Qld only	250g/100L water	In all cases apply to young, actively growing weeds, ensuring thorough coverage.
	Annual & Perennial Pigweed, Artichoke Thistle, Bathurst Burr, Billygoat Weed, Blue Snake Weed, Burr Medic, Clockweed*, Fleabanes, Galvanised Burr, Hemlock, Hoary Cress*, Kyalinga Weed (Whisker Grass), Knobweed, Milky Cotton Bushes, Parthenium Weed, Patersons Curse, Saffron Thistle, Star Burr, Thornapple, Variegated Thistle*		250g/100L water	
	Rubber Vine		125g/10L water	Apply to freshly cut stump.
Conservation Tillage - Direct Drilling, Surface Sowing or Fallow Maintenance	Charlock, Mustards, Shepherd's Purse, Saffron, Slender, Spear and Variegated Thistles, Turnip Weed, Wild Radish, Wild Turnip	All States	400g to 1.3kg	Apply to actively growing young plants before sowing. Observe plant back periods given in the table on this leaflet.
	Clover, Sorrel		900g plus 280-400mL Kamba® 500	Apply to actively growing plants in autumn. Do not sow pasture seed for at least 30 days after application.
Pastures - Spray/Graze Techniques				*PRECAUTION: An increased quantity of poisonous plants may be eaten by stock using Spray-Graze eg. Caltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's Curse.
	Amsinckia, Thistles, Capeweed, Doublegee, Mustard, Paterson's Curse, Wild Turnip, Wild Radish, Docks, Geranium, Erodium	SA only	400g	Apply from 6 weeks after opening rains in autumn until the end of August. Seven days after spraying stock paddock at 4-5 times normal rate, preferably with sheep. Maintain this level of grazing for 6 weeks or until pasture shows signs of over grazing. Then return to normal stocking levels. Use high stocking rates in following spring to prevent weeds from flowering. Repeat treatments may be required for 2-3 years for complete control.
	Annual Thistles, Capeweed, Doublegee, Mustards, Paterson's Curse, Turnip, Saffron Thistle, Spear Thistle	Tas, Vic only		
	Amsinckia, Docks (seedling only), Capeweed, Doublegee, Mustard, Wild Radish, Wild Turnip, Paterson's Curse, Annual Thistles	WA only	500g	
	Spear Thistle, Saffron Thistle		900g	Apply to Saffron Thistle at the end of September when plants are running up to flower. Sub-clovers may be damaged at this rate and use is not recommended for all medic pastures.
	Melons		1.3kg +1% oil	Heavy stocking on young plants sprayed with 650g/ha provides effective control.
	Docks	Vic only	900g	Apply in September only and follow other recommendations above.

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Pastures - Spray/Graze Techniques (cont.)	Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherd's Purse, Saffron, Slender, Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	NSW only	250 to 900g	Spray actively growing 6-8 week old weeds. Introduce stock 7-10 days after spraying, preferably sheep (cattle are less effective). Stocking rate should be at least 5 times heavier than normal until weeds have been reduced, but before survival of desirable pasture species is threatened. Lucerne and Medics may be damaged and should be grazed short before spraying. Other legumes may be affected.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: PASTURE, CEREAL CROPS - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15 APRIL TO 15 SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WEED TABLE:

NOTE: Where weeds are to be sprayed in a CROP or PASTURE use only the rates given for the particular crop or situation indicated under the Directions For Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed.

The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those states where a rate or range of rates is indicated for the particular weeds listed.

WEED TABLE:

Weeds Controlled	Application Rate per Hectare							Critical Comments
	Crop						Pasture -	
	Vic	NSW	SA	Qld	Tas	WA	NSW, SA, Qld, Tas only	
Amaranthus spp.	-	400 to 900g	-	700g	-	-	-	Spray young plants.
Apple of Peru	-	400 to 900g	-	700g	-	-	-	Spray young plants. Susceptible when young.
Bathurst Burr	-	600 to 900g	-	700g	-	-	600 to 900g NOT SA	Spray seedlings only.
Blackberry Nightshade	-	400 to 900g	-	700g	-	-	-	
California Burr	-	600 to 900g	-	700g	-	-	600 to 900g NOT SA	Spray seedlings only.
Cape Tulip	-	-	-	-	-	500g to 1kg	-	Low rate for cormils only
Capeweed	1.1kg	-	1.3kg	-	1.1kg	-	1.3 to 2.1kg	Spray seedlings to rosette stage.
Caltrop	-	600g to 1.3kg	-	700g	-	-	-	Moderately susceptible.
Charlock	400 to 600g	400 to 900g	400g	-	1.1kg	-	600 to 900g	Spray at rosette stage.
Clover	-	900g	-	-	-	-	-	
Common Ice Plant	-	-	900g	-	-	-	-	
Docks	1.1kg	-	1.1kg	1.1kg	1.1kg	900g	2.5kg SA only	Spray at multiple leaf stage. -effective only on seedlings.
Fat Hen	-	400g to 1.3kg	-	700g	1.1kg	-	-	Spray pre-flowering.
Fumitory - Red	-	-	1.3kg	-	-	-	-	
Fumitory - White	600g	-	400g	-	-	-	-	Spray at multiple leaf stage.
Hexham Scent/Melilotus	1.1kg	-	900g	1.1kg	-	-	900g to 1.3kg	Spray multiple leaf stage before seeding.
Hoary Cress	700g to 1.1kg	900g to 1.3kg	1.1kg	1.1kg	-	-	1.1 to 1.3kg	Spray rosettes and pre-flowering.
Hogweed/Wireweed	1.1kg	-	-	1.1kg	-	-	-	Spray at multiple leaf stage (Vic.). Spray at seedling and young plant stage (Qld.)
Horehound	-	-	1.1kg	-	-	-	1.8 to 2.5kg SA ONLY	Spray seedlings.
Khaki Weed	-	-	-	-	-	-	900g to 1.8kg NOT SA	Spray seedlings only.
Lincoln Weed	-	-	1.3kg	-	-	-	-	Spray early rosettes.
London Rocket	-	-	-	-	-	600g	-	
Lupins	-	600g to 1.3kg	-	-	-	-	-	
Mexican Poppy	-	-	-	1.1kg	-	-	-	Spray seedlings - plants become

Weeds Controlled	Application Rate per Hectare							Critical Comments
	Crop						Pasture -	
	Vic	NSW	SA	Qld	Tas	WA	NSW, SA, Qld, Tas only	
Mintweed	-	900g	-	700g	-	-	-	most resistant with age.
Mustards	250 to 400g	400 to 900kg	400g 1.1kg	700g	-	600g	400 to 900g	Spray seedlings - resistant in later stages.
New Zealand Spinach	-	900 to 1.3kg	-	-	-	-	-	Spray at 2-4 leaf up to rosette stage.
Noogoora Burr	-	600 to 900g	-	700g	-	-	600 to 900g NOT SA	
Paterson's Curse	-	900g to 1.3kg	-	1.1kg	-	1kg	1.3 to 1.8kg	Spray seedlings only.
Potato Weed	-	400 to 900g	-	700g	-	-	-	Spray rosettes or before plants have 10 leaves. Later stages harder to kill.
Rapeseed	-	600 to 1.3kg	-	-	-	-	-	
Rough Poppy	-	900g	-	-	-	-	-	
Safflower	-	400 to 900g	-	-	-	-	-	
Shepherd's Purse	-	900g to 1.3kg	-	-	1.1kg	-	600 to 900g	
Skeleton Weed	1.1kg	900g to 1.3kg	1.1kg	-	-	-	1.1 to 1.8 kg	Spray young rosettes.
Sorrel	1.1kg	1.3kg	1.1kg	-	-	-	-	Spray rosettes before aerial growth commences.
Speedwell-Ivy Leaf	-	-	900g	-	-	-	-	Only moderately susceptible.
Spiny Emex	-	-	-	1.1kg	-	-	-	
Stinkwort	-	600g to 1.1kg	-	-	-	-	-	Only young plants are susceptible.
Storkbill/Erodium	-	-	-	-	1.1kg	-	1.3 to 2.5kg	
Sunflower (seedlings)	1.1kg	400g to 1.1kg	-	700g	-	-	-	Spray seedlings to young rosettes.
Thistle - Californian	-	-	-	-	2.1kg	-	2.5 to 2.9kg	
- Saffron	900g	400g to 1.3kg	1.1kg	1.1kg	800g	900g	900g to 1.3kg	Repeated application may be necessary. (NSW, Tas only)
- Slender/Shore	-	600g to 1.3kg	-	-	1.1kg	-	900g	Low rate only sufficient to control weeds in crops at rosette stage when sprayed early.
- Soldier	1.1kg	-	-	-	-	-	900g to 1.3kg NOT NSW, Tas	Suppression only.
- Spear	400g	-	-	-	1.1kg	-	900g to 1.3kg	Spray young rosette.
- Star	-	-	-	-	-	-	1.3 to 1.9kg SA ONLY	Spray young rosettes.
- Variegated	-	400g to 1.3kg	-	700g	1.1kg	-	900g to 1.3kg	Use higher rate as flower stalk appears.
Thornapple	-	600 to 900g	-	-	-	-	1.3 to 1.9kg NOT SA	Spray at rosette stage.
Turnip Weed/Rapistrum	-	400 to 900g	-	400g	-	600g	400 to 900g	Spray seedlings only.
Wards Weed	-	-	900g	-	-	-	-	
Wild Cabbage	1.1kg	-	-	-	-	-	-	
Wild Poppy	400g	-	-	-	-	-	800g to 1.3kg	Spray multiple leaves.
Wild Radish	1.1kg	1.1 to 1.3kg	1.1kg	700g	1.1kg	600g	600 to 900g	Spray rosettes.
Wild Turnip	250 to 400g	400 to 900g	250g	-	1.1kg	500g	400 to 900g	Spray up to young rosette stage.
Vetches/Tares	1.1kg	-	900g	-	-	-	-	Spray 2-4 leaf up to rosette stage.

IMPORTANT

WHEN APPLIED TO DRY SOILS AT LEAST 15mm (1/2inch) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

% In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for Wheat, Barley and Triticale.

In Queensland, planting of Canola/Rapeseed, Chickpeas and Safflower must be delayed for at least 14 days following rainfall of at least 15mm.

@ In Central Queensland, when using 1l/ha or less of Baton, the plant back period for Sorghum and Sunflower is 1 day irrespective of rainfall.

+ In W.A. the plant back period for Lupins at all rates is 28 days.

Crop	Plant back days for Nufarm Baton		
	Rates		
	Up to 400g	400 to 900g	900g to 1.3kg
Balansa Clover	7	7	10
Barley %	1	1	3
Chickpeas #	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins +	7	14	21
Medic	7	7	10
Narbon Beans	7	7	10
Navybean	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10
Phalaris	7	7	10
Canola/Rapeseed #	14	21	28
Rice	7	7	14
Safflower #	7	14	21
Sorghum @	3	7	10
Soybean	14	14	21
Sub-Clover	7	7	10
Sunflower @	7	10	14
Triticale %	1	3	7
Vetch	7	7	10
Wheat %	1	3	7
White Clover	7	7	10

GENERAL INSTRUCTIONS

Before opening, carefully read Directions For Use, Precautionary Statements, Safety Directions and First Aid Instructions.

MIXING

Mix product only with water. Just add the recommended water to your spray tank then remove the water soluble packet from outer bag, add the appropriate amount into the spray tank, agitate for three to five minutes. This product will make a stable water mixture, it is non-flammable. Adding oil, wetting agent or other surfactant to the spray may be used to increase effectiveness on weeds, but doing so may reduce selectivity to crops, resulting in crop damage.

APPLICATION INFORMATION

BOOM SPRAYING -Use 30 to 120L/ha of water.

AERIAL SPRAYING -Use 10 to 90L/ha of water.

EQUIPMENT MAINTENANCE AND USAGE:

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been well washed out with hot soapy water or 1% solution of ammonia, followed by several clear water rinses.

COMPATIBILITY

This product may be tank mixed with Banvel*, Flowable Diuron, Glean*, Nuquat®, Propon®, Nu-trazine, Lusta™, Spray•Seed* and Tordon* 50-D.

RESISTANT WEEDS WARNING

Baton Herbicide ("Baton") is a member of the phenoxy group of herbicides. Baton has the disruptors of plant cell growth mode of action. For weed resistance management Baton is a Group I herbicide. Some naturally-occurring weed biotypes resistant to Baton and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Baton or other Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Baton to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds. DO NOT spray cereals if lucerne is present. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

Drift Warning: Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, vegetables, legume crops and pastures, oilseed crops, susceptible trees (e.g. Kurrajongs, Belahs, Eucalypts) and ornamentals.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container, in a dry, cool, well ventilated area away out of direct sunlight. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty outer containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia **13 11 26**).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

Nufarm Australia Limited ("Nufarm") shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions.

NRA Approval No 46314/0902

In case of emergency: Phone (1 800) 03 3498

Ask for shift supervisor. Toll free 24 hours.

® Nuquat and Propon are registered trademarks of Nufarm Australia Limited.

™ Trademark Pending