

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
CREW™

Herbicide

Active constituent: 330 g/L TRIFLURALIN
Solvent: 364g/L LIQUID HYDROCARBON

GROUP D HERBICIDE

A pre-emergence herbicide for the control of annual grasses and certain broadleaf weeds in certain horticultural and agricultural crops as listed in the Directions for Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT.**

Contents: 5 Litres
20 Litres
110 Litres
200 Litres
1000 Litres

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



™ Trademark pending

GENERAL INSTRUCTIONS

STORAGE AND DISPOSAL

Store in the closed, original container in a cool well-ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Refillable containers [1000L label only]: Store in the closed, original container in a cool well-ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. Do not store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container.

Store the original sealed Envirodrum in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Do not tamper with the Micro Matic valve or the security seal. Do not contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow length PVC gloves, face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone 13 11 26). If swallowed DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

"Nufarm Australia Limited ('Nufarm') shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."

NRA Approval No. 54454/0602

In case of emergency: Phone 1800 033 498

Ask for shift supervisor. Toll free 24 hours.

DrumMuster logo (5L, 20L, 200L labels only)

Envirodrum logo (110L only)

Batch No:

Date of Manufacture:

[Barcode]

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
CREW™

Herbicide

Active constituent: 330 g/L TRIFLURALIN
Solvent: 364g/L LIQUID HYDROCARBON

GROUP D HERBICIDE

A pre-emergence herbicide for the control of annual grasses and certain broadleaf weeds in certain horticultural and agricultural crops as listed in the Directions for Use table.

READ THOROUGHLY BEFORE USING.

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001

NRA Approval No. 54454/0602

™ Trademark pending



DIRECTIONS FOR USE:

1. FIELD CROPS

Situation & Crop	Weeds Controlled	State	Rate L/ha/soil type			Critical Comments
			Light	Medium	Heavy	
Chickpea	Annual Ryegrass, Paradoxa Grass (Canary Grass), Wireweed (Hogweed), Black Pigweed. Suppression of Climbing Buckwheat (Black Bindweed), Wild Oats.	Qld only	1.8-2.4	1.8-2.4	1.8-2.4	Use 1.8L/ha when applying immediately prior to sowing. Use 2.4L/ha when applying to dry soil before the planting rain. Refer to incorporation table 5, 6 or 12 for suitable method of incorporation.
	Annual Ryegrass, Wireweed (Hogweed), Deadnettlles	Vic only	1.2L plus 1.6L Avadex Xtra			Incorporate as per recommendations for wheat, barley and triticale.
	Red & White Fumitory, Rough Poppy, Wireweed, Annual Ryegrass, Barley Grass, Canary Grass, Sand Fescue, Suppression of Deadnettle, Speedwell, Three Cornered Jack, Yellow Burrweed, Brome Grass, Cereal Oats and Surface Wild Oats	SA only	1.8	1.8	1.8	Apply to level seedbed 0 to 4 weeks before sowing. Incorporate as per Instructions Table 5, 6, 12 or 13.
	Amaranthus, Annual Ryegrass, Barnyard Grass, Caltrop, Crab Grass, Paradoxa Grass (Canary Grass), Pigweed, Wild Oats, Winter Grass, Wireweed, Suppression of Fumitory	NSW, ACT only	1.7-2.2	2.2	2.5	Apply from 4 weeks up to just prior to sowing. Refer Incorporation Table 3, 4, 5, 6 or 12 for suitable method of incorporation.
Adzuki Beans, Cowpeas, Lablab, Mung Beans, Borlotti Beans, Red Kidney Beans		NSW, ACT, Qld only				
Faba Beans	Annual Ryegrass, Barley Grass, Capeweed, Corn Gromwell (Sheepweed), Fumitories, Geranium, Ivy Leaf Speedwell, Mustards, Turnips, Wireweed. Suppression of Brome Grass, Soursob and Wild Oats	SA only	1.2L plus 550-830g/ha Nufarm Simazine 900DF (or flowable equivalent)			Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing. Application should not be made to ridged or excessively cloddy soil. For fully reliable results, significant rainfall (20 or 30mm) is necessary within 2-3 weeks of application. Refer to incorporation table 5, 6 or 12 for suitable method of incorporation.
Pigeon Peas	Amaranthus, Barnyard Grass, Canary Grass, Crowsfoot Grass, Pigweed, Spiny Burrgrass, Summer Grass, Wild Oats, Wireweed (Hogweed), Suppression of Yellow Vine (Caltrop), From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass	NSW, ACT only	1.7	2.2	2.5	Apply between 4 weeks and just before sowing. Refer to Incorporation Table 3, 4, 6 or 12 for suitable method of incorporation.
Lentils	Annual Phalaris, Annual Ryegrass, Wild Oats, Wireweed	NSW, ACT only	1.2	1.7	1.7	Apply 0 to 4 weeks before sowing. Refer to incorporation table 5, 6 or 12 for suitable method of incorporation.
	Fumitory – Red and White, Rough Poppy, Wireweed, Barley Grass, Canary Grass, Annual Ryegrass, Sand Fescue	SA only	1.8	1.8	1.8	Apply 0 to 4 weeks before sowing. Sow lentils at a depth of 5cm. Refer to incorporation table 5, 6 or 12 for suitable method of incorporation.

1. FIELD CROPS continued

Situation & Crop	Weeds Controlled	State	Rate L/ha/soil type			Critical Comments
			Light	Medium	Heavy	
Vetch	Annual Ryegrass, Wireweed, soil surface Wild Oats, Suppression of Brome Grass, Rough Poppy, Speedwell, Three Cornered Jack, Yellow Burr Weed, Deadnettle, Sand Fescue, Silver Grasses	SA only	2.4	2.4	2.4	Apply to level seedbed 0 to 4 weeks before sowing. Refer to Table 6, 12 or 13 for method of incorporation.
Cotton	Annual Ryegrass, Barnyard Grass, Canary Grass, Caltrop (Bullhead/Yellow Vine), Crab Grass, Mossman River Grass (Innocent Weed), Pigweed, Redroot (Amaranthus), Redshank (Prince of Wales Feather), Summer Grass, Wild Oats, Winter Grass, Wireweed (Hogweed), Black Pigweed (Qld only), From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	Qld, NSW, WA only	1.7	2.5	3.4	Spray between 6 weeks and just before sowing takes place. Refer Table 1 or 2 for method of incorporation.
Cotton Fallows		NSW only	Apply one of the following rates: FEB/MAR/APR: 4.9L/ha MAY/JUNE: 4.6L/ha JULY: 4.3L/ha AUGUST: 4L/ha			Apply 90 to 100L/ha of water per hectare for ground application. Aerial application – 25-50L/ha water rate. Incorporate (refer incorporation table 1, 2 or 6) as soon as possible after application, as delay may cause inferior weed control.
Legume Seed Crop Establishment - Annual Medics - Clover (Berseem, Red, Strawberry, Sub and White) - Lucerne		NSW, ACT, SA, WA, Vic, Tas only	1.7	1.7	2.5	Autumn Sowing – Apply from 4 weeks to 7 days before sowing takes place. Refer Table 6 for method of incorporation.
			1.7	2.5	2.5	Spring Sowing – Apply between 4 weeks and 3 days before sowing takes place. Refer Table 6 for method of incorporation. In both cases seedling disease, cold weather, excessive moisture, high salt concentrations and drought could weaken crop seedlings and damage could occur from the use of this product. Temporary crop suppression could result.
Linseed		NSW, ACT, SA, WA, Vic only	1.7	2.2	2.5	Spray 2-4 weeks before sowing. Sowing depth should be 1.3-2.5cm. Deeper sowing may result in some stand reduction. Refer Table 6 for method of incorporation.
Peanuts		WA, Qld, NSW, ACT only				Spray between 4 weeks and just before sowing takes place. Refer Table 3, 4, 5, 12 or 13 for suitable method of incorporation.
Peas		All States				Spray between 4 weeks and just before sowing takes place. Refer Table 6 or 12 for method of incorporation.
Rapeseed (Canola) Safflower						
Sugar cane		Qld, NSW only	3.4	3.8	4.3	Apply to plant cane after emergence to "out of hand" stage. Apply to ratoon cane immediately after harvest. Refer Tables 3, 7, 10 or 11 for suitable method of incorporation. Use high rate on heavier soil types. Add Nutrazine 900DF for broadleaf weed control.
Sunflowers		All States	1.7	2.2	2.5	Spray between 4 weeks and just before sowing takes place. Refer Table 3, 4, 5, 12 or 13 for method of incorporation.
Lupins						Spray between 4 weeks and just before sowing takes place. Refer Table 6 or 12 for method of incorporation.
	Annual Grasses and Broadleaf Weeds	NSW, ACT, Vic, Qld only	1.2L plus 1.7kg/ha of Nufarm Simazine 900DF (or flowable equivalent)			Use a low volume boom applying 50-100 litres spray mixture per hectare. Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing the crop.

1. FIELD CROPS continued

Situation & Crop	Weeds Controlled	State	Rate L/ha/soil type			Critical Comments
			Light	Medium	Heavy	
Lupins continued	Annual Grasses and Broadleaf Weeds continued	NSW, ACT, Vic, Qld only	1.2L plus 1.7kg/ha of Nufarm Simazine 900DF (or flowable equivalent)			Do not apply to a ridged soil. Refer to Table 5, 6, 12 or 13 for incorporation method.
	Capeweed, Turnip, Radish, DoubleGee and Suppression of Annual Ryegrass and Wild Oats	WA only	1.8L plus 550-830g/ha of Nufarm Simazine 900DF (or flowable equivalent)			Rate for Yellow Sands. Refer to incorporation table 6, 12 or 13 for incorporation method.
	As above plus Suppression of Brome Grass		1.8L plus 1.1kg/ha of Nufarm Simazine 900DF (or flowable equivalent)			Rate for all other soil types. Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing. Application should not be made to ridged or excessively cloddy soil. For Simazine to be effective sufficient rainfall (20-30mm) to wet the soil through the weed root zone is necessary within 2-3 weeks of application. Results with Simazine can be variable if seasonal conditions are dry prior to sowing and Lupins are sown into dry or low moisture seed beds. Refer to incorporation table 6 or 12 for suitable method of incorporation.
	Capeweed, Doublegee, Wild Radish, Wild Turnip plus suppression of Annual Ryegrass, Wild Oats and Brome Grass		1.8L plus 1.1kg/ha of Nufarm Diuron 900DF (or flowable equivalent)			DO NOT use on white or grey sands as severe crop damage may result. Use tank mix of Diuron & CREW where Annual Ryegrass is present. Apply at pre-sowing stage. Refer to incorporation table 13 for incorporation method.
	Red & White Fumitory, Rough Poppy, Wireweed, Barley Grass, Canary Grass, Annual Ryegrass, Sand Fescue, suppression of Deadnettle, Speedwell, Three Cornered Jack, Yellow Burr Weed, Brome Grass, Cereal Oats, soil surface Wild Oats	SA only	1.8-2.4	1.8-2.4	1.8-2.4	Spray between 4 weeks and just before sowing takes place. Refer Table 6 or 12 for method of incorporation.
	Above weeds plus Capeweed, Common Fumitory, Geranium, Indian Hedge, Mustard, Sheepweed, Shepherd's Purse, Toadrush, Turnips, suppression of Ice Plant, Soursob		1.8-2.4L plus 1.1-2.2kg/ha of Nufarm Simazine 900DF (or flowable equivalent)			Use a low volume boom applying 70-100 litres spray mixture per hectare. Apply to bare moist soil and incorporate to a depth of 5cm just prior to sowing the crop. Do not apply to a ridged soil. Refer to incorporation table 6 or 12 for suitable method of incorporation.
Tobacco	Summer Grass, Crowsfoot Grass, Red Natal Grass, Love Grass, Button Grass, Rhodes Grass, Pigweed	Qld only	1.4	1.7		Apply to soil 3-4 weeks prior to transplanting. The longer period to be used for applications made during June & July. Incorporate to a depth of 10cm. Refer to incorporation table 3 or 9 for incorporation method.
	Crowsfoot Grass	NSW, ACT only	1.4	1.7	1.7	Apply to light sandy soil 10 to 21 days before transplanting. Do not incorporate to a depth greater than 6cm. Apply to loam (medium soil) 7 to 21 days before transplanting. DO NOT incorporate to a depth of greater than 6cm. Refer to incorporation table 3 or 9 for incorporation method.

1. FIELD CROPS continued

Situation & Crop	Weeds Controlled	State	Rate L/ha/soil type			Critical Comments
			Light	Medium	Heavy	
Wheat, Barley & Triticale 1. Pre-Sowing Only 2. Pre-Sowing & Post Sowing (self-mulching soils)	Annual Ryegrass, Wireweed (Hogweed), Phalaris spp.	NSW, ACT, WA, Vic only	1.2	1.2	1.2	Apply 0-28 days before sowing. Sowing depth should be at least 5cm. Use cover harrows behind combine or airseeder. Ground should be left flat. DO NOT use pre-sowing on self-mulching soils as damage may occur from wheel tracking and poor control of wild oats. Refer Table 6 or 12 for method of incorporation.
	Fumitory	WA only				
	Canary Grass	Vic only				
	As above, except for Fumitory	All states				
Wheat	Annual Ryegrass, Paradoxa Grass (Canary Grass), Wild Oats, Wireweed (Hogweed)	Qld only	1.2	1.2	1.2	On non self-mulching soils apply 0-28 days before sowing. Sowing depth should be at least 5cm. Use cover harrows behind combine. Ground should be left flat. On self-mulching soils, as above except apply 1-4 weeks before sowing to prevent crop damage. Refer Table 6 or 12 for method of incorporation.
Barley						Apply to self-mulching and non self-mulching soils from 0-28 days before sowing. Sowing depth should be at least 5cm. Use cover harrows behind combine. Ground should be left flat. Refer Table 6 or 12 for suitable method of incorporation.
Wheat, Triticale, Rye	Annual Ryegrass, Red & White Fumitory, Phalaris spp., Wireweed, suppression of Deadnettle, Rough Poppy, Yellow Burr Weed	SA only	1.2	1.2	1.2	Apply 0-28 days before sowing. Sowing depth should be at least 5cm. Use cover harrows behind combine. Ground should be left flat. DO NOT use pre-sowing on self-mulching soils as damage may occur from wheel tracking and poor control of wild oats. Refer Table 6 or 12 for method of incorporation.
Barley						As above for SA plus Sand Fescue and suppression of Brome Grass
Wheat & Triticale only	Annual Phalaris	NSW, ACT only	1.2L plus 20g/ha Lusta			If possible, spray and incorporate into the soil in one operation. Delay may cause inferior weed control. Refer to incorporation table 12 or 13 for suitable method of incorporation.

2. VEGETABLES, ORCHARDS AND VINEYARDS

Situation & Crop	Weeds Controlled	State	Rate L/ha/soil type			Critical Comments
			Light	Medium	Heavy	
Transplants Only Broccoli, Cabbage, Cauliflowers, Tomatoes	Annual Ryegrass, Barnyard Grass, Canary Grass, Caltrop (Bullhead/Yellow Vine), Crab Grass, Mossman River Grass (Innocent Weed), Pigweed, Redroot (Amaranthus), Redshank (Prince of Wales Feather), Summer Grass, Wild Oats, Winter Grass, Wireweed (Hogweed) From seed only: Columbus Grass, Guinea Grass, Johnson Grass, Liverseed Grass (Urochloa)	All States	1.7	2.5	3.4	Spray between 4 weeks and just before sowing takes place. Refer Table 3, 4, 5 or 11 for suitable method of incorporation.
Direct Seeded Only Broccoli, Brussels Sprouts, Cabbage						
Cauliflower		Vic, Qld only				
Carrots		All States				
Chicory		Vic only				
Green Beans, Navy Beans, Soybeans		All States	1.7	2.2	2.8	
Orchards and Vineyards		Qld, SA, WA, Vic, Tas NSW, ACT only	2	2.5	3.8	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

THIS PRODUCT MUST BE INCORPORATED INTO THE SOIL WITHIN 6 HOURS OF APPLICATION. HIGH APPLICATION VOLUMES (>200L/HA) GIVE OPTIMUM RESULTS.

INCORPORATION TABLE

1. Prior to furrowing out: - 2 workings at an angle required using Offset OR Tandem disc harrows.
2. After furrowing out: - 2 workings required using Go-Devil discs OR Lilliston cultivators set at 10 cm depth.
3. Rotary Hoe: - 1 working required at 5-7.5cm depth. Sugar Cane: - 7.5-13cm depth.
4. Offset OR Tandem Disc Harrows:- (preferably with spiked harrows in tandem) 2 workings at an angle required at 7.5-15cm depth at 6.5-10km per hour.
5. Heavy Diamond OR Stump Jump Harrows:- (weighted 20-30kg per section) at 10-13 km/h speed. Then cross work with offset or tandem disc harrows set to 7.5-15cm depth at speed 6.5-10km per hour.
6. Weighted Heavy Diamond OR Stump Jump Harrows: (weigh with 20-30kg per section) at 10-13km/hr. Cross work with combine at 5-7.5cm depth at speed of 10-13km/hr.
7. Disc Ratoon Cultivator: - 2 workings needed with discs and cultivator set at 7.5-13cm depth.
8. Offset or Tandem Disc Harrows:- set at 7.5-15cm depth. A second discing is required working in opposite direction with discs set to throw treated soil into tree or vine row.
9. Rotary Hoe: - 1 working needed at 5-10cm depth.
10. Offset Discs (Bumpers): - 2 workings needed at depth of 7.5-13cm.
11. Overhead irrigation of 25-50mm or rainfall equivalent (sugar cane and vegetable crops only)
12. Incorporation by sowing on suitably prepared seedbed with heavy diamond harrows trailing or as a separate operation.
13. Post-sowing/Pre-emergence: Use heavy diamond harrows cross working at right angles to the direction of sowing. Do not attempt this method of incorporation on poorly prepared, clumpy or cloddy soils. Reduced or unsatisfactory weed control may occur if heavy weed seed burdens exist and previous tillage or cultivation places these seeds at depth in the soil profile or away from the CREW herbicide.

MIXING

This product is an emulsion which mixes readily with water. Add the recommended amount to the spray tank during filling operation.

APPLICATION

Apply 70-450L of water/ha (broadcast basis) dependent on soil type. More consistent results occur when CREW is applied in higher water rates. Under hot conditions or where possible spray and incorporate into the soil in one operation. Delay may cause inferior weed control. Use properly calibrated standard low pressure (170-340 kilopascal) boomtype sprayer with fan tips.

CONDITIONS FOR BEST RESULTS

This product must be thoroughly incorporated as recommended. Soil should be well worked and free of weeds at time of application. Product effectiveness may be reduced by inadequate incorporation, high organic matter, excess clods, crop or trash residues, stones or other foreign matter and in areas of unnaturally high weed seed population such as header tracks or livestock rest areas. CREW is volatile and disappears from exposed surfaces. Loss is hastened by high temperatures, winds or warm moist soil. Under these conditions apply and incorporate as soon as possible.

WILD OATS

Germinating wild oat seeds lying on soil surface will be controlled. Therefore, specific wild oat control is only possible with shallow cultivation. Poor control will occur on self mulching soils and all soil types where deep cultivation is practiced.

COMPATIBILITY

CREW may be mixed with: Avadex Xtra, Nutrazine 900DF, Nutrazine Flowable, Simazine 900 DF, Simazine Flowable, Diuron 900 DF, Diuron Flowable, Nugran, Logran*, Lusta, Glean*, Nuquat 250, Sprayseed, Nufarm Glyphosate 360, Weedmaster® Duo, Credit® (glyphosate)/Bonus, Glyphosate CT, Tillmaster, Sandoban*, Amicide® 625, Baton®, Surpass® 300, Chlorpyrifos, Dimethoate, Maldison, Endosulfan.

EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible. Otherwise, spray tanks, pumps, lines and nozzles should be thoroughly rinsed several times with clean water following application. Spraymate Tank & Equipment Cleaner is suitable for this purpose and will also remove CREW stains.

RESISTANT WEEDS WARNING

GROUP D HERBICIDE

Nufarm CREW Herbicide ("CREW") is a member of the Dinitroanilines group of herbicides. CREW has the inhibitors of tubulin formation mode of action. For weed resistance management CREW is a Group D herbicide. Some naturally-occurring weed biotypes resistant to CREW and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by CREW or other Group D herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of CREW to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT use in high winds.

DO NOT exceed rates specified, to avoid crop damage.

DO NOT plant sensitive grasses such as oats, sorghum, millets, phalaris spp., ryegrass or wheat for 12 months following the use of this product except where wheat follows wheat or other winter crops.

DO NOT plant oilseed poppies when a detectable residue of trifluralin is present in the soil. Levels as low as 0.02ppm may interact with other unfavourable factors (moisture, stress, disease etc.) to reduce poppy growth and vigour.

DO NOT apply to orchards and vineyards after first flush of growth or when residues can lodge on or in fruit.

Reduced germination of wheat and barley may occur due to combination of following circumstances and the use of this product:

- Short coleoptile cultivars
- Use of seed dressings (except Vitavax)
- Shallow or uneven seedling depth.

Drift Warning: DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool well-ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Refillable containers [1000L only]: Store in the closed, original container in a cool well-ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. Do not store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container.

Store the original sealed Envirodrum in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Do not tamper with the Micro Matic valve or the security seal. Do not contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow length PVC gloves, face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone 13 11 26). If swallowed DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

"Nufarm Australia Limited ('Nufarm') shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."

NRA Approval No. 54454/0602

® Amicide, Baton, Surpass are registered trademarks of Nufarm Australia Limited.

™ Trademark pending.

* Other registered trademarks.

In case of emergency: Phone 1800 033 498

Ask for shift supervisor. Toll free 24 hours.