

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**KENSO**  
agcare

**Abroholos 600**  
Selective Herbicide

ACTIVE CONSTITUENT : 600g/L 2,4-D present as the  
ETHYL HEXYL ESTER

**GROUP I HERBICIDE**


  
**KENSO**  
agcare

A specially formulated low volatile herbicide for selective control of various weeds in crops, pastures and non-agricultural areas as per the Directions for Use Table

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**CONTENTS:**  
20 litres  
APVMA Approval No:  
59709/20/0805

  
9 328666 000952

12/05  
Abroholos 600 20L

Kenso Corporation (M) Sdn Bhd Kirkland Corner H/177 Old Cleveland Rd. Coorparoo 4151 Phone: (07) 3847 4288

**GENERAL INSTRUCTIONS**

Before opening, carefully read Directions For Use, Precautionary Statements, Safety Directions and First Aid Instructions.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilizers. Triple or preferably pressure rinses containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**110L Mini Bulk Returnable Container (110L only)**

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the drum with water or any foreign matter.

After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drum remains the property of Kenso Corporation (M) Sdn Bhd.

**Refillable containers (1000L only):**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**SAFETY DIRECTIONS**

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 13 11 26).

**MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS).

**CONDITIONS OF SALE**

"Kenso Corporation (M) Sdn. Bhd." ("Kenso") shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Kenso's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Kenso has any authority to add to or alter these conditions.

In a Transport Emergency Dial <b>000</b> Police or Fire Brigade
---

Batch No:

Date of Manufacture:

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Kenso Agcare**  
**Abroholos 600**

**Selective Herbicide**

**ACTIVE CONSTITUENT: 600 g/L 2,4-D present as the  
ETHYL HEXYL ESTER**

**GROUP I HERBICIDE**

A specially formulated low volatile herbicide for selective control of various weeds in crops, pastures and non-agricultural areas as per the Directions for Use Table.

**This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals.**

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT



**Kenso Corporation (M) Sdn Bhd**  
Kirkland Corner H/177 Old Cleveland  
Rd. Coorparoo 4151  
Phone: 07 3847 4288

**DIRECTIONS FOR USE:**

**RESTRAINTS:** AVOID spraying if rain is likely within 6 hours or if strong winds prevail. DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

**1. FIELD CROPS**

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
<b>Wheat, Barley</b>	Refer to Weed Table	Vic only	240-900mL	CROP STAGES:ALL CEREALS Apply at tillered to boot stages. (WA, Vic only).  Apply after when the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller. (NSW, ACT only). Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling of the head at the top of the main stem. (Qld only). Apply from completion of tillering to early jointing stage. (SA, Tas only).
		SA only	260-900mL	
		QLD, NSW, ACT only	460-900mL	
		Tas only	700-900mL	
		WA only	900mL	
<b>Triticale</b>		SA, Qld, NSW, ACT only	460-900mL	
<b>Cereal Rye</b>		Vic only	240-900mL	
		NSW, Qld, ACT only	460-900mL	
<b>Sugar cane</b>		Vic only	240-900mL	
		NSW, Qld, ACT only	460-900mL	
<b>Stubble/ Fallow Spray Prior to Direct Drilling or Sowing. Winter Cereals, Grain legumes (Peanuts Qld only), Canola</b>		Qld only	1.3-2.7L	Post-emergence
		All States	240-900mL/ha	Observe the plant back periods given in the table in this leaflet. Must be tank mixed with a knockdown herbicide such as Roundup <sup>^</sup> Max, Credit <sup>®</sup> plus Bonus <sup>®</sup> , Ken-up 450 CT, Glyphosate CT, Paraken 250 or Speedy 250. Select appropriate rate from the weed table. For skeleton weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.
<b>Harvest Aid or Salvage Spray</b>	Broadleaf Weeds	All States	1.9L	Apply after dough stage of crop. Interval between application and effectiveness is 10-20 days. For desiccation of green matter, estimate harvest date and apply spray approximately 14 days earlier. Rain between spraying and actual harvest can negate results. NB. Where thistles are tall and branching above the crop, spraying can turn the branches down into the crop, presenting more stalks to cause header comb blockages. Spraying may increase seed contamination of harvest by accelerating maturity. Do not use with undersown legumes that have not set seed.
<b>-Winter Cereals</b>	Refer to Weed Table			
<b>Potatoes Pre-harvest Preparation</b>	Broadleaf Weeds such as Clover, Variegated Thistle & Cruciferous weeds	Vic, Tas only	1.3-2.7L	Apply approximately 4 to 5 weeks before harvest after the potato haulms have dried off. Use the highest rate where weeds are more than 30cm in height. For boom spraying apply at least 100 litres of spray mixture per hectare. If grasses such as Rye grass and winter grass are also present add Amitrole T Herbicide.

**2. PASTURES, NON-AGRICULTURAL, INDUSTRIAL**

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
<b>Improved Pasture containing Clovers</b>	Refer to Weed Table	NSW, Tas, SA, Qld, ACT only	460-900mL	Clover must be well covered by the grass or extensive damage may result.
<b>Pastures – non legumes, Rights of way &amp; Industrial</b>		NSW, Tas, SA, WA, Qld, ACT only	900mL-5.3L	Control of most perennial weeds, but due to the rooting habits of most species control may take a number of years. Damage may result to legumes in pasture.
		Vic only	900mL -7.5L 80-700mL	Boom spray Spot spraying
<b>Pastures – Direct Drilling or Surface Sowing</b>	Charlock, Clover, Medics, Mustards, Paterson's Curse, Saffron, Slender, variegated and Spear Thistles, Turnip weed, wild Radish, wild Turnip	NSW, ACT only	900mL-1.7L (Aerial application)	Apply to young, actively growing weeds. Sowing: Do not sow pasture seed for at least 21 days after application. If soil moisture is dry, delay sowing for at least 30 days.
	As Above plus: Capeweed, Wireweed, Storksbill/Erodium, Flatweed, Horehound (seedlings), Skeleton Weed, Nodding or Star Thistles		900mL – 1.3L (Ground Application)	
	St. John's Wort		3.7-5.3L (Aerial or Ground)	
	All of Above plus grasses		As above plus Propon or Roundup^ Max or Credit® plus Bonus® or Glyphosate CT or Weedmaster® Duo	

**3. SPOT SPRAYING**

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
<b>Spot Spraying (All Situations)</b>	Refer to Weed Table	All States	1/100 <sup>th</sup> of rate on Weed Table per 10L water per 100m <sup>2</sup>	Apply through Knapsack. Thorough wetting of weed is essential.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.  
IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15<sup>TH</sup> APRIL TO 15<sup>TH</sup> SEPTEMBER UNLES OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.**

**WITHHOLDING PERIOD (WHP): PASTURES, CEREALS CROPS – DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

**HARVEST WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.**

**GENERAL INSTRUCTIONS**

Before opening, carefully read Directions For Use, Precautionary Statements, Safety Directions and First Aid Instructions.

**APPLICATION INFORMATION**

This product may be used in either high and low volume sprays. Just pour into water and stir.

BOOM SPRAYING – Use 30-100 litres water per hectare.

AERIAL SPRAYING – Use 10-90 litres water per hectare.

NOTE: Refer to the Department of Agriculture/ Primary Industries in your state for the current restricted spraying areas.

**EQUIPMENT MAINTENANCE AND USAGE**

Keep the spray unit for herbicides only if possible. Otherwise wash out the unit with hot soapy water followed by several clear water rinses. Do not use wooden spray vats as they can not be cleaned.

Hoses cannot be cleaned and new hoses should be fitted when the unit is to be used for any other purpose.

**COMPATIBILITY**

This product can be tank mixed with 500g/L Dicamba, Ken-Chlor 750, Paraken 250, Propon®, Atrazine 900 WG Herbicide, Credit®/Bonus®, Roundup<sup>^</sup> Max, Roundup<sup>^</sup> CT, Ken-up 450 CT, Ken-Met 600, Ken-Gran 750, Speedy 250\*, Weedmaster Duo.

**WEED TABLE**

NOTE: Where weeds are to be sprayed in a CROP or PASTURE, use only the rates given for the crop in the table below. In most cases this will give control, however some hard to kill weeds may only be suppressed. E.g. Rumex spp. (docks) and Polygonum spp. (wireweed, climbing buckwheat) are killed to ground level only.

**APPLICATION RATE PER HECTARE**

Weeds Controlled	CROP						PASTURES –NON LEGUMES		
	Vic	SA	Tas	NSW, ACT	Qld	WA	Vic	NSW, Tas, SA, Qld, WA, ACT only	Critical Comments
Amaranthus spp	-	-	-	900mL	-	-	-	-	
Angled Onion	-	-	-	-	-	-	3.7L	0.9-1.9L	Spray when buds forming or early flowering
Apple of Sodom	-	-	-	-	-	-	-	3.3-3.7L	
Bathurst Burr	-	-	-	900mL	-	-	1.9-3.7L	1.9-3.7L	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Black Knapweed	-	-	-	-	-	-	3.7L	-	Spray before flowering. DO NOT cultivate these infestations
Buffalo Burr	-	-	-	-	-	-	-	900mL – 1.3L (not Qld & WA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
California Burr	-	-	-	900mL	-	-	1.9-3.7L	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Caltrop	-	-	-	700-900mL	-	-	1.9-3.7L	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Cape Tulip	-	-	-	-	-	1.3L	3.7L	1.9-3.7L	Spray before flowering.
Capeweed	900mL	900mL	900mL	600-900mL	-	-	-	2.8-3.7L	Spray up to rosette stage
Charlock	460mL	460mL	900mL	460mL	-	-	-	900mL	Spray up to rosette stage.
Clover	-	-	-	700-900mL	-	-	-	-	
Colocynth	-	-	-	-	-	-	3.7L	-	Spray at seedling stage only

Deadnettle	-	-	-	900mL	-	-	-	-	
Devil's Claw	-	-	-	900mL	-	-	1.5L	-	Spray prior to pods forming.
Dock	900mL	900mL	-	-	900mL	900mL	-	1.9-2.8L	Spray at rosette stage to kill top growth only
Fat Hen	-	-	-	460-900mL	-	-	-	-	
Flatweed	-	-	-	900mL	-	-	-	2.8-3.7L	
Fumitory (red)	-	900mL	-	900mL	-	-	-	2.8-3.7L	Spray up to rosette stage
Fumitory (white)	900mL	460mL	-	900mL	-	-	-	2.8-3.7L	Spray up to rosette stage
Galvanised Burr	-	-	-	-	-	-	5.3L	5.3L (not Qld & WA)	Spray from seedling to pre-flowering.
Goosefoots	-	-	-	900mL	-	-	-	-	
Hard Head or Russian Knapweed	-	-	-	-	-	-	3.7-5.9L	-	Spray before flowering
Hogweed, Wireweed	900mL	900mL	-	900mL	900mL	-	-	1.3-1.9L (not SA)	Spray up to rosette stage
Hoary Cress, Whiteweed	-	900mL	900mL	900mL	-	-	1.9-3.7L	1.9-2.4L	Spray from late rosette to pre-flowering.
Horehound (seedlings)	-	900mL	-	-	-	950mL	-	1.9-3.7L	Late Autumn to early Spring
Ironweed, Corn Gromwell	-	-	-	900mL	-	-	-	1.3-1.9L	
Khaki Weed	-	-	-	-	-	-	-	900mL-1.3L (Not SA)	Spray young seedlings only
Lincoln Weed	-	900mL	-	-	-	-	-	-	Autumn spray before sowing improves control.
London Rocket	-	-	-	-	-	650mL	-	1.8-2.8L (WA only)	
Lupins	900mL	-	-	460mL - 900mL	-	-	-	-	Spray up to rosette stage.
Melilotus/ Hexham Scent	900mL	900mL	-	-	900mL	-	-	1.3-1.9L	Spray up to rosette stage.
Melons – camel, paddy	-	-	-	460mL - 900mL	-	-	-	-	
Mustards	370mL	260-900mL	900mL	460mL - 900mL	700mL	700mL	3.7L	1.9-2.8L	Spray up to rosette stage.
Mexican Poppy	-	-	-	900mL	-	950mL	-	900mL-1.3L (1.3-1.7L WA only)	Spray rosette stage and before flowering.
Mintweed	-	-	-	900mL	700mL	-	-	900mL-1.3L	Spray active seedlings only
Muskweed	900mL	-	-	-	-	-	-	-	Spray up to rosette stage.
New Zealand Spinach	-	-	-	900mL	-	-	-	-	
Noogoora Burr	-	-	-	900mL	-	-	1.9-3.7L	1.9-3.7L	Spray seedling to pre-flowering
Nut Grass	-	-	-	-	-	-	3.7-5.9L	-	Spray within 4 weeks of foliage emergence, repeat spray necessary.
Paterson's Curse	-	-	-	900mL	-	950mL	1.9-3.7L	900mL-1.9L(1.3-1.7L WA only)	Spray seedling to rosette stage

Poppy Wild	460mL	-	-	-	-	-	-	2.4-3.3L	Spray up to rosette stage.
Ragwort	-	-	-	-	-	-	3.7L	3.7L	Spray at rosette to cabbage stage.
Rapeseed	900mL	-	-	460mL - 900mL	-	-	-	-	Spray up to rosette stage.
Rapistrum	-	-	-	-	-	650mL	-	950mL (WA only)	
Rough Poppy	-	460mL	-	460mL - 900mL	-	-	-	900mL	Spray young seedlings only.
St. John's Wort	-	-	-	-	-	-	3.7- 5.9L	3.7-5.3L	Spray before flowering. Spray before plants are 40cm high.
Safflower	-	-	-	460mL - 900mL	-	-	-	-	
Sand Mustard/ Sand Rocket	-	-	-	-	-	-	3.7L	-	Spray before flowering
Shepherd's Purse	-	-	-	900mL	-	-	-	-	
Silver Leaf Nightshade	-	-	-	-	-	-	3.7L	-	Spray at flowering. Fallowland: controls top growth only
Skeleton Weed	-	900mL	-	900mL	-	-	3.7L	1.3-1.9L	Spray rosettes before aerial growth commences.
Stingless Nettle (Deadnettle)	-	900mL	900mL	-	-	-	-	2.4-2.8L	
Stinging Nettle	900mL	-	-	-	-	-	-	-	Spray up to rosette stage
Stinkwort	-	-	-	900mL	-	-	1.9- 3.7L	1.9-3.7L	Spray younger plants. Use higher rate as plants mature.
Storksbill/ Erodium	-	-	-	900mL	-	-	-	-	
Sunflower Seedlings	900mL	-	-	460mL - 900mL	700mL	-	-	-	Spray multiple leaves.
Thistles:	-	-	-	-	-	-	-	-	
-Golden	-	-	-	-	-	-	3.7L	3.7L	Spray at rosette stage
-Nodding	-	-	-	-	-	-	3.7L	1.3-1.9L	Spray rosette stage to pre-flowering
-Saffron	700mL	900mL	-	460mL - 900mL	900mL	900mL	900mL- 1.9L	900mL- 2.8L	Spray up to rosette stage.
-Sheep	-	-	-	-	-	950mL	-	950mL- 3.7L (WA only)	
-Slender, Shore	-	-	900mL	900mL	-	-	1.9- 3.7L	900mL- 3.7L	Spray at rosette stage.
-Soldier	-	-	-	-	-	-	3.7L	-	Spray at rosette stage
-Spear	900mL	-	900mL	-	-	-	900mL- 2.8L	1.3-2.4L	Spray at seedling to rosette stage. Use higher rate as plants mature (pastures)
-Stemless	-	-	-	-	-	-	3.7L	2.8-3.7L	Spray rosette stage to flowering.
-St. Barnaby's	-	-	-	-	-	-	-	1.3-1.9L	
- Star	-	-	-	900mL	-	-	1.9- 3.7L	-	Spray seedling to rosette stage. Use higher rate as plants mature.

-Variegated	-	-	900mL	460mL- 900mL	700mL	-	900mL- 2.8L	900mL- 3.7L	Spray at rosette stage. Can cause stock poisoning.
Thornapple	-	-	-	460mL - 900mL	-	-	3.7L	900mL- 1.9L	Spray at seedling stage
Tree Hogweed	900mL	-	-	-	-	-	-	-	Spray up to rosette stage
Turnip Weed	-	460mL	-	460mL - 900mL	460mL	700mL	-	900mL	Spray seedlings only
Vetches/Tares	900mL	700mL	900mL	-	-	-	-	-	
Wards Weed	-	460mL	-	-	-	-	-	-	Spray at seedling stage
Wild cabbage	900mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Wild Garlic	-	-	-	-	-	-	7.5L	-	Suppresses aerial growth only
Wild Mignonette	-	-	-	-	-	950mL	3.7L	-	Spray at rosette stage
Wild Mustard	-	-	-	-	-	650mL	-	1.8-2.8L (WA only)	
Wild Radish	900mL	900mL	900mL	460mL - 900mL	900mL	650mL	-	900mL (950mL WA only)	Spray up to rosette stage
Wild Sage	-	-	-	-	-	-	-	2.8-3.7L	
Wild Teasel	-	-	-	-	-	-	1.9- 3.7L	-	Spray at rosette stage. Use higher rate as plants mature.
Wild Turnip	240mL	260mL	900mL	460mL - 900mL	-	450mL	-	900mL (950mL WA only)	Spray up to rosette stage.

## Plant back days for Kenso Agcare Abroholos 600 Selective Herbicide

Crop	Rates		
	Up to 580mL/ha	580mL-1.15L/ha	1.15-1.8L/ha
Balansa Clover	7	7	10
Barley%	1	1	3
Chickpeas#	7	14	21
Cotton	10	14	21
Faba beans	7	7	10
Field peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins +	7	14	21
Medics	7	7	10
Narbon beans	7	7	10
Navybean	10	10	14
Oats	3	3	7
Perennial ryegrass	7	7	10
Persian clover	7	7	10
Phalaris	7	7	10
Canola/ rapeseed #	14	21	28
Rice	7	7	14
Safflower#	7	14	21
sorghum@	3	7	10
Soybean	14	14	21
Sub-clover	7	7	10
Sunflower@	7	10	14
Triticale%	1	3	7

<b>Vetch</b>	7	7	10
<b>Wheat%</b>	1	3	7
<b>White Clover</b>	7	7	10

**IMPORTANT:WHEN APPLIED TO DRY SOILS AT LEAST 15MM (½inch) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.**

**NOTES:**

% In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for wheat, barley and Triticale.

# In Queensland, planting of canola/rapeseed, chickpeas and safflower must be delayed for at least 14 days following rainfall of at least 15mm.

@ In Central Queensland, when using 830mL /ha or less of Abroholos 600, the Plant Back Period for sorghum and sunflower is 1 day irrespective of rainfall.

+ In WA the Plant Back Period for Lupins at all rates is 28 days.

**RESISTANT WEEDS WARNING**

Kenso Agcare Abroholos 600 Selective Herbicide is a member of the Phenoxy group of herbicides. Abroholos 600 has the Disruptors of plant cell growth mode of action. For weed resistance management Abroholos 600 is a Group I herbicide. Some naturally-occurring weed biotypes resistant to Abroholos 600 and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Abroholos 600 or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use. Kenso Corporation accepts no liability for any losses that may result from the failure of Abroholos 600 to control resistant weeds.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

**Legume Tolerance:** If clovers are present, care should be taken to ensure that they are reached the 3-4 leaf stage before spraying. Rates above 460mL of this product per hectare will destroy most clovers, whilst Lucerne and medic are susceptible at any strength.

**Drift Warning:** DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. In particular, avoid spray drift and vapour movement onto susceptible crops such as: cotton, tobacco, tomatoes, flowers, vegetables, vines, fruit trees, legume crops and pastures, oilseed crops or other susceptible plants and trees (eg. Kurrajongs, Eucalypts and Belahs).

DO NOT use unless wind speed is more than 3 kilometres per hour and less than 15 kilometres per hour as measured at the application site. DO NOT apply with smaller than coarse to very coarse spray droplets according to the BCPC/ASAE S572 definition for standard nozzles.

**PROTECTION OF LIVESTOCK**

Low hazard to bees. May be applied at any times as recommended in the Directions for Use.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or a place away from children, animals, food, feedstuffs, seed and fertilizers. Triple or preferably pressure rinses containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**110L Mini Bulk Returnable Container (110L only)**

Store the original sealed drum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the drum with water or any foreign matter.

After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drum remains the property of Kenso Corporation (M) Sdn Bhd.

**Refillable containers (1000L only):**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**SAFETY DIRECTIONS**

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 13 11 26).

**MATERIAL SAFETY DATA SHEET**

For further Information refer to the Material Safety Data Sheet (MSDS).

**CONDITIONS OF SALE**

"Kenso Corporation (M) Sdn. Bhd." ("Kenso") shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Kenso's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Kenso has any authority to add to or alter these conditions.

In a Transport Emergency Dial <b>000</b> Police or Fire Brigade
---

Batch No:

Date of Manufacture:



Kenso Corporation (M) Sdn. Bhd.  
Kirkland Corner H/177 Old Cleveland Rd.  
Coorparoo 4151

APVMA Approval No: 59709/0805