

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Access*

Herbicide

ACTIVE CONSTITUENTS: 240 g/L TRICLOPYR present as the butoxyethyl ester
120 g/L PICLORAM present as the isooctyl ester

SOLVENTS: 389 g/L LIQUID HYDROCARBON

GROUP I HERBICIDE

For selective control of a wide range of woody and noxious weeds in commercial and industrial areas, public lands, fence lines and pastures, by basal bark and cut stump applications as specified in the Directions For Use.

Dow AgroSciences Australia Limited A.B.N. 24 003 771 659

20 Rodborough Road FRENCHS FOREST NSW 2086

www.dowagrosciences.com.au

CUSTOMER SERVICE TOLL FREE 1-800 700 096

* Trademark of Dow AgroSciences

DIRECTIONS FOR USE

RESTRAINTS

DO NOT add water to Access Herbicide.

DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result.

DO NOT apply to charcoal coated or wet stems as this can repel the diesel mixture.

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FENCE LINES, FORESTRY, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /60 L DIESEL	CRITICAL COMMENTS
<i>Acacia</i> spp. including: Black wattle Brigalow Hickory wattle Mimosa bush Mulga Sally wattle Silver wattle	Basal bark method: Plants with stems up to 5 cm basal diameter.	1 L	Ensure all stems on multi-stemmed varieties are treated. Delay treatment until all regrowth on bulldozed or ploughed land has had time to emerge to a height of at least one metre. Root suckering wattles will need follow-up treatment.
African boxthorn	Cut Stump method: Plants up to and in excess of basal bark size. Apply immediately after cut is made.		Treat all stems on multi-stem plants.
Algaroba (<i>Prosopis</i> spp.)			
<i>Angophora</i> spp. regrowth			This species may become dormant so should not be treated during winter.
Australian blackthorn			
Bitterbark		Regrowth from roots may occur.	
Broadleaf privet		2 L	
Brush box			Treat single stemmed seedling plants only.
Cacti including: Common pest pear Harrisia cactus Prickly pear Snake cactus Tiger pear Tree pear		1 L	Apply as an overall spray, wetting all areas of the plant to ground level.
Cockspur thorn			Basal bark application only. Species with old, rough bark require more thorough wetting than smooth barked species. Some regrowth may still occur.
Corkwood wattle			
<i>Corymbia</i> spp.			
Ellangowan poison bush			
<i>Eucalyptus</i> spp. regrowth			
False sandalwood			
Green cestrum			
<i>Grevillea</i> spp.			
Groundsel bush			

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FENCE LINES, FORESTRY, PASTURES AND RIGHTS-OF-WAY.					
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /60 L DIESEL	CRITICAL COMMENTS		
Guava	Basal bark method: Plants with stems up to 5 cm basal diameter.	2 L	Regrowth may occur from roots of the main tree.		
Hard milkwood		1 L			
Hawthorn					
Heartleaf poison bush					
Honey locust					
Kiteleaf poison	Cut Stump method: Plants up to and in excess of basal bark size. Apply immediately after cut is made.				
Leucaena					
Limebush					Treat single stem plants only.
Lantana					
Narrow leaf hop bush					
Needlewood					
Oleander					
Parkinsonia					
Punty bush					
Red ash (white myrtle)					
Rubbervine					
Swamp box		2 L	Treat single stemmed seedling plants only.		
Sweet briar		1 L	Do not treat during winter.		
Tea trees					
Turpentine					
Turpentine bush					
Whitewood					
Wild orange	Basal bark treatment only.				
Wild peach					
Wild Rosemary					
Wild tobacco tree					
Wilga	Cut stump treatment only.				
Yellow bells (<i>Tecoma</i> spp.)				Some root suckering may occur.	

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FENCE LINES, FORESTRY, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /60 L DIESEL	CRITICAL COMMENTS
Camphor laurel	Basal bark method: Plants with stems up to 10 cm basal diameter.	1 L	Species that become dormant should not be treated during winter.
Peppercorn tree (<i>Schinus molle</i>)			Basal bark method only.
Prickly acacia			
Sisal hemp			Basal bark method only.
Tree of Heaven	Cut Stump method: Plants up to and in excess of basal bark size. Apply immediately after cut is made.	4 L	Species that become dormant should not be treated during winter.
Willow			Cut stump method only. Complete control may not occur due to the multi-stem growth of plant and difficulty treating all stems.
Chinee apple	Basal bark method: Plants with stems up to 15 cm basal diameter. Cut stump method: Plants up to and in excess of basal bark size. Apply immediately after cut is made.	1 L	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

FOR NATIVE VEGETATION

Use of Access on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

WITHHOLDING PERIOD:

The use pattern of this product is such that no withholding periods are required.

GENERAL INSTRUCTIONS

MIXING

Access herbicide is designed for use with diesel distillate only. It will NOT mix with water.

Quarter fill the spray unit or mixing container with diesel and add the required amount of Access. Add the remaining diesel and shake or agitate thoroughly to mix the contents. Only mix sufficient chemical for each day's work, mixing periodically.

Dilution Table for one part Access in 60 parts diesel:

Amount of Access Herbicide	Volume of diesel added after Access
17 mL	85 mL
255 mL	1.7 L
fill to 1 L	fill to 5 L
fill to 15 L	fill to 100 L

COMPATIBILITY

Access is compatible with diesel and should not be used or diluted with any other chemicals.

GENERAL APPLICATION

Apply only with hand-directed equipment such as a pneumatic sprayer. The use of diesel as a herbicide carrier may affect the rubber seals in some sprayers. To avoid this, use sprayers, which use Viton® seals and fittings.

Spray equipment should be used at low pressures, up to 200 kPa, to avoid splashing.

Ensure all stems on multi-stemmed varieties are treated. Delay treatment until all regrowth on bulldozed or ploughed land has had time to emerge to a height of at least 1 metre.

Species that lose their leaves e.g. sweet briar; or become dormant, e.g. Australian blackthorn or camphor laurel, should not be treated during winter.

Species with old, rough bark require more thorough wetting than the smooth barked species.

DO NOT apply to charcoal coated or wet stems as this can repel the diesel mixture.

Basal Bark Application Method

Liberaly spray the bark around the stem from ground level to 30 cm high, wetting thoroughly to the point of runoff (unless otherwise stated).

Cut Stump Method

Stems should be cut less than 15 cm above ground level.

Immediately apply Access/diesel mixture **liberaly** to the **freshly** cut stump by spraying the cut surface and sides of the stem.

CLEANING SPRAY EQUIPMENT

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

After completion of spraying, use a degreaser such as Caltex Kwik-D-Grease® to remove traces of diesel from the sprayer. Rinse tank and spray through nozzles thoroughly with water to remove degreaser.

After the above, quarter fill the tank with clean water and add a alkali detergent, (e.g. liquid SURF®, OMO®, DRIVE® at 500 mL/100L of water or the powder equivalent at 500 g/100L of water). If using a concentrated laundry detergent use 250 g (or mL)/100 L water.

Do not use chlorine based cleaners.

Shake the water around the sprayer and spray the solution through the nozzles to thoroughly clean the lines and nozzles. Rinse well with clean water to remove detergent.

To clean brushes and container, spray liberally with degreaser. Hose off thoroughly with water and repeat using detergents (see above).

DO NOT use this equipment for any other purpose.

RESISTANT WEEDS WARNING

GROUP I HERBICIDE

Access Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management the product is a Group I Herbicide.

Some naturally occurring weed biotypes resistant to the product and other disrupters of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group I Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Dow AgroSciences accepts no liability for any losses that may result from the failure of Access Herbicide to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Dow AgroSciences representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Treat target weeds only. **DO NOT** allow mixture to contact non-target species.

DO NOT use on land to be cultivated for growing sensitive plants. Legumes, vines, vegetables, cotton, tomatoes, ornamentals, coniferous species and many other broadleaf plants are highly susceptible to Access during both growing and dormant periods.

Picloram, one of the active ingredients in this product, can remain active in the soil for extended periods depending on soil type, rainfall, temperature, humidity, soil moisture and soil organic matter.

DO NOT move soil, which has been sprayed, to areas where desirable plants are to be grown.

DO NOT apply when the treated soil may be washed to areas growing, or to be planted with desirable plants. DO NOT apply close to, or on, areas containing roots of desirable plants, or sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted to, susceptible crops or plants.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

Alongside waterways treat only noxious weeds and poisonous plants.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

DO NOT store near food, feedstuffs, fertilisers or seed. Triple rinse empty containers with diesel before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site.

The method of disposal of the container depends on the container type. Read the STORAGE AND DISPOSAL instructions on the label that is attached to the container.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand or clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

SAFETY DIRECTIONS

- Harmful if swallowed.
- Will irritate the eyes, nose, throat and skin.
- Repeated exposure may cause allergic disorders.
- Avoid contact with eyes and skin.
- Avoid inhaling vapour or spray mist.
- When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrists, a washable hat and elbow-length neoprene gloves and a face shield or goggles.
- If product on skin, immediately wash area with soap and water.
- If product in eyes, wash it out immediately with water.
- After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.
- After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

- If poisoning occurs, contact a doctor or Poisons Information Centre. (Phone: Australia 13 11 26)
- If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet for Access Herbicide, which is available from Dow AgroSciences on request. Call Customer Service Toll Free on 1-800 700 096 or visit www.agrosciences.com.au

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions for use. No warranty of merchantability or fitness for a particular purpose express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by Dow AgroSciences, or under abnormal conditions.

APVMA Approval No. 46640/0906

* Trademark of Dow AgroSciences

® Registered Trademarks

IN A TRANSPORT
EMERGENCY ONLY
DIAL 000
FOR POLICE OR
FIRE BRIGADE

EMERGENCY RESPONSE
(ALL HOURS)
RING FROM ANYWHERE
IN AUSTRALIA
1-800 033 882
(LOCAL CALL FEE ONLY)

