

# CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## AGCHLOR 750 WG HERBICIDE

ACTIVE CONSTITUENT: 750 g/kg CHLORSULFURON

GROUP	<b>B</b>	HERBICIDE
-------	----------	-----------

A selective herbicide for the control of Annual (Wimmera) Ryegrass and certain broadleaved weeds in Wheat, Barley, Oats, Cereal Rye and Triticale.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS: 500g, 1kg, 2kg, 5kg, 10kg

### ALLFIRE ENTERPRISES PTY LTD

1/61 Dowd St  
Welspool, WA, 6106  
Tel: 08 92585755  
ACN 097 970 280

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## **SAFETY DIRECTIONS**

Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Tel: 131126).

## **MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet which is available from the supplier.

## **TERMS OF SALE:**

Allfire Enterprises Pty Ltd will not be held liable for any loss, injury or damage, indirect or consequential, arising from the sale, supply, use or application of this product. The product is not to be used for any purpose or in any way contrary to label instructions.

APVMA Approval Number: 56878/0803

Batch No:

Date of Manufacture:

# CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## AGCHLOR 750 WG HERBICIDE

ACTIVE CONSTITUENT: 750 g/kg CHLORSULFURON

GROUP	<b>B</b>	HERBICIDE
-------	----------	-----------

A selective herbicide for the control of Annual (Wimmera) Ryegrass and certain broadleaved weeds in Wheat, Barley, Oats, Cereal Rye and Triticale.

This leaflet is part of the label

NRA APPROVAL NO: 56878/0803

### **ALLFIRE ENTERPRISES PTY LTD**

1/61 Dowd St  
Welspool, WA, 6106  
Tel: 08 92585755  
ACN 097 970 280

**DIRECTIONS FOR USE**

**RESTRAINTS:**

DO NOT spray emerged crops if rain is expected within four hours.

After mixing in the tank, spray within 48 hours if this product is used by itself, or within 24 hours if mixed with another product.

DO NOT apply to plants suffering stress.

**METHOD OF USE - PRE-SOWING INCORPORATED BY SOWING**

**Annual Ryegrass**

Crop/Situation	Weeds Controlled	State(s)	Rate g/ha			Critical Comments
			Soil Type			
			Light to Medium Soils	Heavy Soils		
			Soil pH			
			Less than 7	7.0 - 8.5	8.5 or less	
Wheat and Triticale only	Annual (Wimmera) Ryegrass <i>Lolium rigidum</i>	NSW, ACT, Vic, SA, WA only	20	15 or 20*	20	* Use the higher rate when paddock history suggests a high weed population can be expected. NOTE: Refer to General Instructions for optimum application timing and conditions.

Crop/Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
Wheat and Triticale only	African Turnip Weed <i>Sisymbrium thellungii</i>	NSW, ACT and Qld only	20	
	Amsinckia/Yellow Burrweed <i>Amsinckia spp.</i>	NSW, ACT, Vic, SA, WA only	15	
	Annual Phalaris <i>Phalaris paradoxa</i> , <i>Phalaris minor</i>	NSW, ACT only	20 + 1 L/ha Trifluralin	
	Barley grass <i>Hordeum leporinum</i>	NSW, ACT and Tas only		
	Silvergrass <i>Vulpia spp.</i>	Tas only		
	Ball Mustard <i>Neslia puniculata</i>	SA only	15	
	Black Bindweed/Climbing Buckwheat <i>Fallopia convolvulus</i>	Qld only	20	Apply to dry soil before the sowing rain. Mechanical incorporation before the sowing rains is not necessary.
	Brome grass <i>Bromus spp.</i> (Suppression only)	NSW, ACT, Vic, SA, WA, Tas only		Gives suppression only if populations are 20 plants/m <sup>2</sup> or less.
	Cape Tulip <i>Homeria spp.</i>	WA only	20	
	Capeweed <i>Arctotheca calendula</i>	NSW, ACT, Vic, SA, WA, Tas only		On acid soils pH 5.5 or less, this product will give a shorter period of control in wet years.
	Charlock <i>Sinapis arvensis</i>	Vic, SA, Tas only	15	
	Common Iceplant <i>Mesembryanthemum crystallinum</i>	SA only		
	Corn Gromwell/Sheepweed/ White Ironweed <i>Buglossoides arvensis</i>	Qld, NSW, ACT, Vic, SA, WA only	20	
	Deadnettle <i>Lamium amplexicaule</i>	All States	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.
	Docks <i>Rumex spp.</i>	NSW, ACT, Vic, SA, WA, Tas only	20	
	Fat-hen <i>Chenopodium album</i>	NSW, ACT, Tas only		
Fumitory <i>Fumaria spp.</i>	NSW, ACT, Vic, SA, WA, Tas only	15 or 20	Use the higher rate when paddock history suggests a high weed population can be expected.	

Crop/Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
Wheat and Triticale only	Guildford grass/Onion grass <i>Romulea rosea</i>	WA only	15	Apply to dry soil before the sowing rain. Mechanical incorporation before the sowing rains is not necessary.
	Indian Hedge Mustard <i>Sisymbrium orientale</i>	All States		
	King Island Melilot <i>Melilotus indicus</i>	Vic, SA only		
	Lincoln weed <i>Diplotaxis tenuifolia</i>	SA only		
	Loosestrife	Vic only		
	Mintweed <i>Salvia reflexa</i>	Qld and NSW, ACT only	20	
	Mouse-ear Chickweed <i>Cerastium spp.</i>	NSW, ACT, Vic, SA, WA, Tas only	15	
	New Zealand Spinach <i>Tetragonia tetragonoides</i>	Qld only	20	
	Paradoxa grass <i>Phalaris paradoxa</i>	Nth NSW, ACT (Soil pH >7.5) and Qld only		
	Paterson's Curse/Salvation Jane <i>Echium plantagineum</i>	NSW, ACT, Vic, SA, WA, Tas only	15	
	Pimpernels <i>Anagallis arvensis</i>	NSW, ACT, Vic, SA, Tas only		
	Prickly Lettuce/Whip Thistle <i>Lactuca serriola</i>	Vic, SA only	20	Use the higher rate when paddock history suggests a high weed population can be expected.
	Rough Poppy <i>Papaver hybridum</i>	NSW, ACT, SA, WA, Tas only	15 or 20	
	Saffron Thistle <i>Carthamus lanatus</i> (Suppression only)	Qld, NSW, ACT, Vic, SA, Tas only	20	
	Saltbush <i>Atriplex muelleri</i>	Qld and NSW, ACT only		Use the higher rate when paddock history suggests a high weed population can be expected.
	Shepherd's Purse <i>Capsella bursa-pastoris</i>	NSW, ACT, Vic, SA, WA, Tas only	15 or 20	
	Slender Celery <i>Apium leptophyllum</i>	Qld and NSW, ACT only	20	
	Slender Thistle <i>Carduus tenuiflorus</i>	Tas only		
	Soursob <i>Oxalis pes-caprae</i>	NSW, ACT, Vic, SA only	15	Apply only to soils of pH 7.5 or above. Apply after majority of Soursobs have emerged and leave soil undisturbed for 1-4 weeks prior to cultivation or sowing. The most effective and reliable control is achieved with early post-emergence applications (EPE) after crop and weed emergence.
	Spear Thistle <i>Cirsium vulgare</i>	Tas only	20	Use the higher rate when paddock history suggests a high weed population can be expected.
	Stemless Thistle <i>Onopordum acaulon</i>	SA only	15 or 20	
	Storksbill/Wild Geranium <i>Erodium spp.</i>	Vic, SA, WA, Tas only	15	
	Three-Cornered Jack(s)/Doublegee/ Spiny Emex <i>Emex australis</i>	NSW, ACT, Vic, SA, WA only	20	
	Tree Hogweed <i>Polygonum patulum</i>	Vic, SA only		
	Turnip weed <i>Rapistrum rugosum</i>	Qld and SA only	15	
	Wireweed/Hogweed <i>Polygonum aviculare</i>	All States	15 or 20	
	Wild Turnip <i>Brassica tournefortii</i>	NSW, ACT, Vic, SA, WA and Tas only	15	

## METHOD OF USE - POST CROP AND WEED EMERGENCE

### Annual Ryegrass

Crop/ Situation	Weeds Controlled	State(s)	Rate g/ha			Critical Comments
			Soil Type			
			Light to Medium Soils	Heavy Soils		
			Soil pH			
Less than 7	7.0 - 8.5	8.5 or less				
Wheat, Barley, Oats, Triticale and Cereal Rye only	Annual (Wimmera) Ryegrass <i>Lolium rigidum</i>	NSW, ACT, Vic, SA, WA only	20 or 25*	15 or 20*	20 or 25*	* Use the higher rate under heavy weed pressure. Apply no later than the 3 leaf stage of Annual Ryegrass. * Application of This product to Annual Ryegrass 2 leaf or greater with water volumes less than 50 L/ha may result in reduced efficacy.

Crop/Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
Wheat, Barley, Oats, Triticale and Cereal Rye only	African Turnip weed <i>Sisymbrium thellungii</i>	NSW, ACT and Qld only	20	Apply at cotyledon to 4 leaf stage.
	Amsinckia/Yellow Burrweed <i>Amsinckia spp.</i>	NSW, ACT, Vic, SA, WA only	15	
	Ball Mustard <i>Neslia punculata</i>	SA only		
	Bifora/Carrot weed <i>Cotula australis</i>		25	
	Black Bindweed/Climbing Buckwheat <i>Fallopia convolvulus</i>	Qld and NSW, ACT only	20	Apply at cotyledon to 2 leaf stage of weed.
	Cape Tulip <i>Homeria spp.</i>	WA only		
	Charlock <i>Sinapis arvensis</i>	NSW, ACT, Vic, SA, Tas only	15	
	Corn Gromwell/Sheepweed/ White Ironweed <i>Buglossoides arvensis</i>	NSW, ACT, Vic, SA, WA only	20	Apply at cotyledon to 2 leaf stage. If applied at a later stage only suppression will occur.
	Deadnettle <i>Lamium amplexicaule</i>	Qld, NSW, ACT, Vic, SA, Tas only	15 or 20	Use the higher rate under heavy weed pressure.
	Docks <i>Rumex spp.</i>	Vic, SA, WA, Tas only	15	
	Fat-hen <i>Chenopodium album</i>	NSW, ACT, Tas only	20	
	Fumitory, Denseflower <i>Fumaria densiflora</i>	NSW, ACT, Vic, SA, WA, Tas only		Apply at cotyledon to 2 leaf stage.
	Guildford grass/Onion grass <i>Romulea rosea</i>	WA only	15	
	Hoary Cress <i>Cardaria draba</i>	Vic, SA, Tas only	20	Apply when plants are fully emerged.
	Lincoln weed <i>Diploaxis tenuifolia</i>	SA only	20	
	Matricaria <i>Matricaria matricarioides</i>	WA, Tas only		
	Mintweed <i>Salvia reflexa</i>	Qld and NSW, ACT only		Apply at cotyledon to 4 leaf stage.
	Mouse-ear Chickweed <i>Cerastium spp.</i>	NSW, ACT, Vic, SA, WA, Tas only	15	
	Mustards <i>Sisymbrium spp.</i>	All States		
	New Zealand Spinach <i>Tetragonia tetragonoides</i>	Qld only	20	
	Paterson's Curse/Salvation Jane <i>Echium plantagineum</i>	NSW, ACT, Vic, SA, WA, Tas only	15	
	Pimpernels <i>Anagallis arvensis</i>	NSW, ACT, Vic, SA, Tas only		
	Prickly Lettuce/Whip Thistle <i>Lactuca serriola</i>	Vic, Tas only	20	
	Rough Poppy <i>Papaver hybridum</i>	NSW, ACT, SA, WA, Tas only		
	Saltbush <i>Atriplex muelleri</i>	Qld and NSW, ACT only		Apply at cotyledon to 4 leaf stage.

Crop/Situation	Weeds Controlled	State(s)	Rate g/ha	Critical Comments
Wheat, Barley, Oats, Triticale and Cereal Rye only	Shepherd's Purse <i>Capsella bursa-pastoris</i>	NSW, ACT, Vic, SA, WA, Tas only	20	
	Slender Celery <i>Apium leptophyllum</i>	Qld and NSW, ACT only		Apply at cotyledon to 4 leaf stage.
	Soursob <i>Oxalis pes-caprae</i>	NSW, ACT, Vic, SA, WA only		Apply when the majority of Soursobs have emerged.
	Spear Thistle <i>Cirsium vulgare</i>	Tas only		
	Stagger weed <i>Stachys arvensis</i>	Qld, NSW, ACT, WA, Tas only		
	Stemless Thistle <i>Onopordum acaulon</i>	Vic only	25	
	Storksbill/Wild Geranium <i>Erodium spp.</i>	Vic, SA, WA, Tas only	15	
	Tree Hogweed <i>Polygonum patulum</i>	Vic only	20	
	Turnip Weed <i>Rapistrum rugosum</i>	Qld, NSW, ACT, SA only	15	
	Wild Radish <i>Raphanus raphanistrum</i>	All States	15 or 20	Use the higher rate under heavy weed pressure. A follow-up spray with a suitable herbicide may be necessary to control subsequent germinations.
	Wild Turnip <i>Brassica tournefortii</i>	NSW, ACT, Vic, SA, WA, Tas only	15	
	Wireweed/Hogweed <i>Polygonum aviculare</i>	All States	20	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED**

#### GENERAL INSTRUCTIONS

This product is a selective herbicide designed to control certain weeds in wheat, triticale, barley, oats and cereal rye.

This product is suitable as a pre-sowing treatment for wheat and triticale, and as a post-sowing treatment for wheat, triticale, barley, oats and cereal rye. When used on emerged weeds, the product is absorbed by foliage and green stems and moves into the root system.

Prior to using this product, careful consideration should be given to soil pH. As soil pH increases, rate of breakdown decreases. This product should not be used on soil pH 8.6 or higher as extended soil residual activity could adversely affect crop rotation options beyond normal intervals.

This product should not be used on soils with a pH of 8.6 or higher as soil residual activity could adversely affect following crops and crop rotation intervals may be extended beyond normal intervals.

Crops other than wheat, barley, oats, triticale and cereal rye can be extremely sensitive to low concentrations of this product in the soil. See Crop Rotation Recommendations.

Best weed control is obtained when rainfall or sprinkler irrigation wets the soil to a depth of 5 to 7.5 cm within 4 weeks of application.

#### Pre-sowing incorporated by sowing:

WA only - Avoid applying to dry sandy soils as rapid leaching may occur with early season rains.

SA only - Before using rates greater than 15 g/ha on light to medium soils pH 7 to 8.5, seek further advice.

Conventionally Sown Crops - on soils less than 7, apply a spray just before sowing or in conjunction with the sowing operation. On soils of pH of 7 or greater it is not critical to time the spray just before sowing. Spray onto a non-ridged surface free of large clods. Use low profile 10 cm combine points for sowing. Sow at speeds of 10 kph or greater. Use light covering harrows at sowing. If applied to

dry soil and sowing is to be delayed, incorporate immediately after spraying to prevent loss by wind erosion.

Direct Drilled Crops - apply tank mixed with either Spray.Seed<sup>1</sup> or glyphosate in accordance with manufacturer's label recommendations.

#### **Post Crop and Weed Emergence:**

Where treatment is delayed or where weeds are not actively growing due to adverse conditions results may be slow to appear and weeds may be only stunted or suppressed.

Wheat, Triticale and Cereal Rye - apply after crop emergence when weeds are small and actively growing (**Annual Ryegrass no more than 3 leaves, Broadleaved weeds no more than 5 cm in height or diameter** (for Black Bindweed refer to specific recommendations)).

Barley and Oats - apply between 2 leaf stage of crop (3 leaf stage - SA only) and early tillering when weeds are small and actively growing (**Annual Ryegrass no more than 3 leaves, Broadleaved weeds no more than 5 cm in height or diameter** (for Black Bindweed refer to specific recommendations)).

#### **Resistant Weeds Warning**

<b>GROUP</b>	<b>B</b>	<b>HERBICIDE</b>
--------------	----------	------------------

Agchlor 750 WG Herbicide is a member of the sulfonylurea group of herbicides. Agchlor 750 WG Herbicide has the inhibitor of the enzyme acetolactate synthase (ALS) mode of action. For weed resistance management, Agchlor 750 WG Herbicide is a Group B herbicide.

Naturally-occurring weed biotypes resistant to Agchlor 750 WG Herbicide and other ALS inhibitor herbicides (Annual Ryegrass and some broadleaf weeds) are known to exist.

They can eventually dominate the weed population if these herbicides are used repeatedly. These weeds will not be controlled by Agchlor 750 WG Herbicide or other ALS inhibitor herbicides.

Annual Ryegrass biotypes resistant to diclofop-methyl and other "grass specific" herbicides are often also resistant to Agchlor 750 WG Herbicide. Before using Agchlor 750 WG Herbicide on a population resistant to "grass specific" herbicides, have a resistance test conducted to ensure that it is still susceptible to Agchlor 750 WG Herbicide.

Since the occurrence of resistant weeds is difficult to detect prior to use, Allfire Enterprises Pty Ltd accepts no liability for any losses that may result from the failure of Agchlor 750 WG Herbicide to control resistant weeds.

To prevent, or at least minimise the risk of resistant weeds occurring, use Agchlor 750 WG Herbicide in tank mixes (if appropriate) and/or rotations with herbicides having different modes of action effective on the same weed species.

Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be taken to prevent seed set of these survivors.

DO NOT make more than one application of an ALS inhibitor herbicide to a crop, either pre-sowing incorporated by sowing or post crop and weed emergence.

If the user suspects that an ALS inhibitor-resistant weed is present, Agchlor 750 WG Herbicide or other ALS inhibitor herbicides should not be used.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

#### **Grazing Advice**

Avoid grazing treated areas within 24 hours of application to optimise weed control.

A nil withholding period is applicable for grazing Agchlor 750 WG Herbicide treated areas (when used as directed on this label).

#### **Crop Safety:**

DO NOT use this product for:

- crops other than cereals.
- cereals irrigated by furrows or flooding.
- winter cereals undersown with legume pasture crops.
- weed control where crops are under stress. Damage can occur where crops are stressed due to conditions such as excessive soil alkalinity or acidity, poor nutrient status, disease, nematode or insect infestation, adverse weather conditions, drought or waterlogging. If crops become stressed after spraying, they may turn yellow or become retarded, but usually they will recover with no reduction in yield.

**Wheat:**

DO NOT use this product for:

- wheat varieties Cranbrook, or Miling.
- the wheat variety Vulcan if on acid soils and under stress conditions caused by waterlogging, frost, aluminium or manganese toxicity; reduced yields may result.
- pre-sowing treatment of weeds in wheat varieties Avocet and Durati (OK for post-emergent use).
- pre-sowing treatment of weeds in wheat variety Banks if soil pH is 5.5 or less (OK for post-emergent use).

**Barley and Oats:**

DO NOT use this product for:

- application before the crop has reached the 2-leaf stage (3-leaf stage in SA).
- Stirling barley.
- barley under waterlogged conditions (yield may be reduced).

The application of other sulfonylurea herbicides following this product is not recommended.

**Crop Rotation Recommendations**

Land previously treated with this product should not be rotated to other crops other than those listed in the following tables.

Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas.

The treated areas may be replanted to any of the specified crops after the interval indicated in the following tables:-

**NB - THE TABLE BELOW APPLIES TO ALL STATES**

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)						
SOIL pH*	0	3	6	9	12	18
6.5 or less	Triticale Wheat	Cereal Rye	Oats	Barley	Subterranean Clover** Faba Beans Field Pea Linseed Lucerne Lupins Medics ** Rapeseed Safflower	Maize Sorghum Soybeans Sunflower

**NB - THE TABLES BELOW APPLY TO Qld, SA, WA & Tas ONLY**

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)						
RAINFALL REQUIREMENT	0	3	9	15	18	22
	Minimum 700 mm					
Soil pH* 6.6-7.5	Triticale Wheat	Cereal Rye	Barley Oats	Japanese Millet Maize Panicum Millet Sorghum Sunflower White French Millet	Cotton Soybeans	Faba Beans Field Pea Linseed Medics** Rapeseed Safflower Subterranean Clover**

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)				
RAINFALL REQUIREMENT	0	15	18	24 months or longer
	Minimum of 700 mm			
Soil pH* 7.6-8.5	Triticale Wheat	Japanese Millet Maize Panicum Millet Sorghum Sunflowers White French Millet	Barley Oats Cereal Rye	Rotate to crops other than Cereals (such as listed above) only if field test strip of the planned rotational crop has been successfully grown through to maturity in the previous season.
8.6 and above	This product is not recommended for use on soils of pH 8.6 and above.			

**NB - THE TABLES BELOW APPLY TO NSW, ACT & Vic ONLY**

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)					
SOIL pH*	0	3	9	22	26
6.6 - 7.5	Triticale Wheat	Cereal Rye	Barley Oats	Subterranean Clover** Faba Beans Field Peas Linseed Lucerne Lupins Medics ** Rapeseed Safflower	Maize Sorghum Soybeans Sunflower

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)			
SOIL pH*	0	18	24 months or longer
7.6 - 8.5			Rotate to crops other than Cereals (such as listed above) only if field test strip of the planned rotational crop has been successfully grown through to maturity in the previous season.
8.6 and above	This product is not recommended for use on soils of pH 8.6 and above.		

\* Soil pH is to be determined by Laboratory analysis using the 1:5 soil:water suspension method.

\*\* Includes natural regeneration of Subterranean clover and Medics.

- Land previously treated with this product should not be rotated to crops other than those listed in the above table.

- Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas.

### Spray Preparation

This product is a water dispersible granule.

1. Fill tank partially with water and engage full agitation.
2. Add the required amount. (N.B. The measuring flask provided is graduated in grams of Agchlor 750 WG Herbicide only. DO NOT use for measuring of other materials.)
3. Top up with water to the required volume.
4. Companion products: If applying this product with another product ensure this product has completely dissolved before adding the companion product.
5. Agchlor 750 WG Herbicide must be kept in suspension at all times by continuous agitation. Where prepared spray mixes have been allowed to stand, thoroughly re-agitate before using.

### Use of Surfactant/Wetting Agent

For post-emergence application, always add a non-ionic surfactant (1000 gac/L) at 100 mL/100 L of final spray volume (0.1% volume/volume).

The use of spraying oils is not recommended.

NOTE: DO NOT add a surfactant/wetting agent when this product is tank mixed with another product which already has a surfactant/wetting agent in the formulation.

### Ground Spraying Equipment

Use a boom spray properly calibrated to a constant speed and rate of delivery to ensure thorough coverage and a uniform spray pattern. Avoid overlapping and shut off spray booms while starting, turning, slowing or stopping as injury to the crop may result. Apply a minimum of 30 L prepared spray/ha.

### Aerial Application

Apply a minimum of 20 L/ha water. Avoid spraying in still conditions and in winds likely to cause drift onto adjacent sensitive crops. Avoid spraying where drift can go onto areas likely to be sown to sensitive crops - see Crop Rotation Recommendations. Turn off spray boom whilst passing over creeks and dams.

### Sprayer Cleanup

It is essential that the sprayer be properly cleaned after using this product to prevent injury to crops other than wheat, triticale, barley, oats, or cereal rye. All traces of Agchlor 750 WG Herbicide should be removed from equipment using the following procedure:

1. Drain tank, then flush tank, boom and hoses with clean water for a minimum of 10 minutes.
2. Fill the tank with clean water then add 300 mL household chlorine bleach (containing 4% chlorine) per 100 L of water. Flush through boom and hoses then allow to sit for 15 minutes with agitation engaged, then drain.
3. Repeat Step 2.
4. Nozzles and screens should be removed and cleaned separately. To remove traces of chlorine bleach, rinse the tank thoroughly with clean water and flush through hoses and boom.

**CAUTION:** DO NOT use chlorine bleach with ammonia. All traces of liquid fertiliser containing ammonia, ammonium nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odour which can cause eye, nose, throat and lung irritation. DO NOT clean equipment in an enclosed area.

### **Compatibility**

This product is compatible with glyphosate and paraquat. This product does not control Wild Oats, however it is compatible with Wild Oat herbicides Avadex BW<sup>1</sup>, Mataven<sup>1</sup>. It is also compatible with Bromoxynil, MCPA (and Bromoxynil/MCPA mixtures), 2,4-D amine and 2,4-D ester, Tigrex<sup>1</sup> and Jaguar<sup>1</sup>. This product is also compatible with trifluralin and the insecticides omethoate, dimethoate deltamethrin and chlorpyrifos.

### **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply or drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

### **STORAGE AND DISPOSAL**

Store in the closed, original container in a well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### **SAFETY DIRECTIONS**

Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.

### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Tel: 131126).

### **MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet which is available from the supplier.

### **TERMS OF SALE**

Allfire Enterprises Pty Ltd will not be held liable for any loss, injury or damage, indirect or consequential, arising from the sale, supply, use or application of this product. The product is not to be used for any purpose or in any way contrary to label instructions.

<sup>1</sup> Not a trademark of Allfire Enterprises Pty Ltd