

**POISON**  
**KEEP OUT REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **ALLFIRE FORCE<sup>TM</sup>**

# **FUNGICIDE**

**ACTIVE CONSTITUENT: 250g/L FLUTRIAFOL**

|              |          |                  |
|--------------|----------|------------------|
| <b>GROUP</b> | <b>C</b> | <b>FUNGICIDE</b> |
|--------------|----------|------------------|

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the Directions for Use Table.

**CONTENTS: 5L, 10L, 20L**

**ALLFIRE ENTERPRISES PTY. LTD**  
**1/61 Dowd St**  
**Welspool, WA, 6106**  
**Tel: 08 92585755**  
**ACN 097 970 280**

<sup>TM</sup> Allfire Enterprises Pty Ltd

## DIRECTIONS FOR USE FOR ALL STATES

### A. APPLIED AS AN IN-FURROW TREATMENT

| CROP   | DISEASE  | RATE PER HECTARE | WHP               | CRITICAL COMMENTS  |
|--------|--|------------------|-------------------|--|
| Canola | Blackleg ( <i>Leptosphaeria maculans</i> )   | 400mL            | 4 weeks (grazing) | Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties.<br>Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400mL of the product on the quantity of fertiliser required to be sown in the furrow per hectare, using the table below as a guide.<br><b>Fertiliser rate per hectare (kg)</b> <b>Volume of product per 100kg of fertiliser (mL)</b><br>60                                      667*<br>70                                      571<br>80                                      500<br>100                                     400<br>120                                     333   |
| Wheat  | Stripe rust ( <i>Puccinia striiformis</i> )  | 200 or 400mL     |                   | For the control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected.<br>Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400mL of the product on the quantity of fertiliser required to be sown in the furrow per hectare, using the table below as a guide.<br><b>Fertiliser rate per hectare (kg)</b> <b>Volume of product per 100kg of fertiliser (mL)</b><br>200mL rate      400mL rate<br>50                                      400                                      800*<br>60                                      333                                      667*<br>70                                      286                                      571<br>80                                      250 <sup>†</sup> 500<br>100                                     200 <sup>†</sup> 400<br>120                                     167 <sup>†</sup> 333 |
|        | Septoria tritici blotch ( <i>S. tritici</i> )<br>Take-All ( <i>Geaumannomyces graminis</i> ) | 400mL            |                   |  |
| Barley | Powdery Mildew ( <i>Erysiphe graminis</i> )<br>Scald ( <i>Rhynchosporium secalis</i> )       | 200 or 400mL     |                   | *Application volumes over 600mL per 100kg of fertiliser may adversely effect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers.<br><sup>†</sup> Where application volume is low, product may be diluted with water to improve coverage.  |

### B. APPLIED AS FOLIAR SPRAY

|        |  |   |                               |  |
|--------|--|---|-------------------------------|--|
| Wheat  | Stripe rust ( <i>Puccinia striiformis</i> )  | 250-500mL<br><b>Ground application:</b><br>Apply in 100-150L water plus 200mL of a 1000g/L non-ionic surfactant per 100L of spray solution. | 7 weeks (Harvest or grazing)  | Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with a high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.                        |
|        | Leaf rust ( <i>P. recondita</i> )<br>Septoria nodorum blotch ( <i>Septoria nodorum</i> ),<br>Septoria tritici blotch ( <i>S. tritici</i> ) | <b>Aerial application</b><br>Apply in a minimum of 15L water plus 200mL a 1000g/L non-ionic surfactant per 100L of spray solution.          |                               | Apply after the flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with a high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate. |
| Barley | Powdery mildew ( <i>Erysiphe graminis</i> )  |   | 10 weeks (harvest or grazing) | Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. In crops with a high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.  |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## **WITHHOLDING PERIODS:**

### **A. FORCE APPLIED IN-FURROW AS A FERTILISER TREATMENT**

CANOLA, WHEAT AND BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.

### **B. FORCE APPLIED AS A FOLIAR SPRAY**

BARLEY – DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION

- DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 10 WEEKS OF TREATMENT.

WHEAT - DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION

- DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 7 WEEKS OF TREATMENT.

## **GENERAL INSTRUCTIONS**

### **Fertiliser application**

Ensure the treated fertiliser is dry before sowing the crop.

Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. DO NOT mix seed with treated fertiliser.

### **Foliar Applications**

Check the crop carefully and regularly. Spray only if there is sufficient time for the rust to develop to a level on the variety in question when it will cause significant yield loss.

### **Fungicide Resistance Warning**

|       |          |           |
|-------|----------|-----------|
| GROUP | <b>C</b> | FUNGICIDE |
|-------|----------|-----------|

Allfire Force Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group C fungicide.

Some naturally occurring individual fungi resistant to Allfire Force Fungicide and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Allfire Force Fungicide or other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Allfire Enterprises Pty Ltd accepts no liability for any losses that may result from the failure of Allfire Force Fungicide to control resistant fungi.

### **Protection of Wildlife, Fish, Crustaceans and Environment**

DO NOT contaminate streams, rivers or waterways with chemical or used containers.

### **Storage and Disposal**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling break, crush, or puncture and bury containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### **Safety Directions**

Harmful if swallowed. Will irritate the eyes. May irritate the nose, throat and skin. Avoid contact with eyes and skin. Avoid inhaling spray mist. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and a face shield or goggles.

If product in eyes, wash out immediately with water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

### **First Aid**

If poisoning occurs contact a doctor or Poisons Information Centre(Tel: 131126)

**Material Safety Data Sheet**

For further information refer to the Material Safety Data Sheet which is available from the supplier.

**NOTICE TO BUYER**

Allfire Enterprises Pty. Ltd. will not be held liable for any loss, injury or damage, indirect or consequential, arising from the sale, supply, use or application of this product. The product is not to be used for any purpose or in any way contrary to label instructions.

APVMA Approval No: 56178/5L/0204  
56178/10L/0204  
56178/20L/0204

Batch No:  
Date of Manufacture: